



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXIII International Grassland Congress

Functional Characterization of Siberian Wild Rye Grass *EsHSP* 16.9 Gene Conferring Diverse Stress Tolerance in Prokaryotic Cells

Ki-won Lee

National Institute of Animal Science, South Korea

Ki-yong Kim

National Institute of Animal Science, South Korea

Hee-chung Ji

National Institute of Animal Science, South Korea

Tae-young Hwang

National Institute of Animal Science, South Korea

Sang-hoon Lee

National Institute of Animal Science, South Korea

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/23/4-1-4/5>

The XXIII International Grassland Congress (Sustainable use of Grassland Resources for Forage Production, Biodiversity and Environmental Protection) took place in New Delhi, India from November 20 through November 24, 2015.

Proceedings Editors: M. M. Roy, D. R. Malaviya, V. K. Yadav, Tejveer Singh, R. P. Sah, D. Vijay, and A. Radhakrishna

Published by Range Management Society of India

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Functional characterization of Siberian wild rye grass EsHSP 16.9 gene conferring diverse stress tolerance in prokaryotic cells

Ki-Won Lee, Ki-Won Lee, Ki-Yong Kim, Hee Chung Ji, Tae Young Hwang, Sang-Hoon Lee

National Institute of Animal Science, Cheonan, Korea

Corresponding author e-mail: kiwon@korea.kr

Keywords: Abiotic stress, Molecular chaperone, Siberian wild rye grass, Small heat shock, Tolerance

Introduction

Siberian wild rye (*Elymus sibiricus* L.) is a perennial, caespitose, and self-pollinating grass indigenous to Northern Asia and also is widely distributed from Northern Europe to Japan. The plant shows strong environmental adaptability with tolerance to drought and cold; thus, it is often used as forage resources (Yan *et al.*, 2007). Environmental stresses caused by global warming are acknowledged to be as a serious issue in agriculture due to reductions of crop productivity (Ahuja *et al.*, 2010). Genetic natural breeding of Siberian wild rye would potentially increase the productivity of forage crops; however, genetic studies on this grass have yet to be conducted. Heat shock proteins (Hsps) are the well characterized stress inducible proteins playing as molecular chaperones in prokaryotes and eukaryotes. We have also identified two differently localized small Hsps: rice chloroplastic and alfalfa mitochondrial Hsps confer tolerance to oxidative and heat stresses in tall fescue and to salinity and arsenic stresses in *E. coli*, tobacco, and tall fescue, respectively (Lee *et al.*, 2012a; Lee *et al.*, 2012b). Here, we cloned the small *Hsp16.9* gene from various heat stress-induced fragments in Siberian wild rye using differentially expressed gene (DEG) analysis. We examined the mRNA expression of *EsHsp16.9*, *in vitro* molecular chaperone activity and *in vivo* stress tolerance by using a prokaryotic system against diverse environmental stresses.

Materials and Methods

Plant materials and growth conditions: Seeds of Siberian wild rye were collected from Ulaanbaatar, Mongolia. Seeds were planted in pots containing Horticulture Nursery medium and grown in a growth chamber maintained at 25°C with a light intensity of 400 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and a 14-h photoperiod.

Isolation of EsHsp16.9: Four-week-old seedlings were subjected to 42°C heat treatments or 25°C non-treatment (control) for 6 h. After treatment, leaves were harvested and frozen with liquid nitrogen. Total RNA was isolated from the leaf tissues of treated and control plants using TRIzol reagent (Qiagen, CA, USA). Using the GeneFishing™ DEG kit (Seegene, South Korea), *EsHsp16.9* was isolated using an annealing control priming-based polymerase chain reaction method, as described by Lee *et al.*, 2012c.

Sequence analyses: The nucleotide and deduced amino acid sequences and the conserved domain were analyzed using NCBI-BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) and Bioedit (version 7.2.3) software. The isoelectric point (pI) and molecular weight (Mw) of *EsHsp16.9* were calculated using the Compute pI/Mw tool in ExPASy (http://web.expasy.org/compute_pi/). Amino acid sequences of small Hsps in *Arabidopsis* and homologs of *EsHsp16.9* from different plant species were retrieved from the GenBank™ database after a homology search using blastp in NCBI-BLAST. Multiple sequence alignment was performed using ClustalX2 and GeneDoc (version 2.7) software. A phylogenetic tree was constructed by the neighbor joining method with 1000 bootstrap replicates by software MEGA6.0 (<http://www.megasoftware.net>).

Abiotic stress treatments to Siberian wild rye plants: Four-week-old Siberian wild rye seedlings were exposed to conditions of drought (withdrawing irrigation), heat (42°C), cold (4°C), salt (NaCl, 300 mM), copper (CuSO₄, 500 μM), cadmium (CdCl₂, 500 μM), arsenate (Na₂HAsO₄·7H₂O, 500 μM), methyl viologen (MV, 1 mM), and hydrogen peroxide (H₂O₂, 10 mM). For heat and cold treatments, seedlings were transferred into a temperature-controlled growth chamber (42 or 4°C). For salt and heavy metal treatments, seedlings were irrigated with water containing indicated concentrations of salt and heavy metals. For MV and H₂O₂ treatments, leaf segments were floated on water containing indicated concentrations of MV and H₂O₂. Stress-treated plants were harvested, frozen in liquid nitrogen and maintained at -80°C until use.

AAB39856), *Medicago truncatula* (Hsp18.2 class I; XP_003619703), *Nicotiana tabacum* (Hsp 3B class I; ADK36668), and *Cicer arietinum* (Hsp8.5 class I; XP_004505085). **(B)** Phylogenetic analysis of EsHsp16.9 with homologs from different plants. The scale bar represents the number of amino acid substitutions per site.




Fig. 2. Expression of EsHsp16.9 transcripts in response to abiotic stressors. Expression of EsHsp16.9 in seedlings treated with heat (a), cold (b), salt (c), drought (d), copper (e), cadmium (f), arsenate (g), methyl viologen (h) and hydrogen peroxide (i) for 0, 1, 6, 12, 24 and 36 h. Three independent northern blot analyses were performed and the band intensity was analyzed using ImageJ. Data are mean \pm SE (n = 3)

Conclusion

Siberian wild rye grass is an important forage crop in northern Asia and has strong adaptability to unfavorable environments. Here, we screened for useful genetic materials from Siberian wild rye by DEG analysis using the gene-fishing technique. Using this approach, we identified a small Hsp 16.9 (EsHsp16.9) which we isolated, successfully cloned, and categorized as falling into the cytosolic class I subfamily. Enhanced expression of *EsHsp16.9* transcripts were also confirmed under various environmental stressors. Small Hsp genes are found in various organisms including plants, act as molecular chaperones to protect cellular proteins against damage from diverse stressors, and have been known to induce stress tolerance. Using a recombinant protein expression system in *E. coli*, we expressed and purified recombinant EsHsp16.9 protein and found that it effectively exhibited molecular chaperone activity to protect thermal aggregation of malate dehydrogenase. We also revealed that *E. coli* cells expressing EsHsp16.9 exhibited better growth under salt,

arsenate, PEG, and heat stressors. Thus, *via* genetic manipulation the *EsHsp16.9* gene has the potential to confer tolerance into crops to multiple environmental stressors.

References

- Ahuja, I., R. C. H. de Vos, A. M. Bones and R. D. Hall. 2010. Plant molecular stress responses face climate change. *Trends Plant Sci.* 15:664-674.
- Lee, K. W., G. J. Choi, K. Y. Kim, H. J. Ji, H. S. Park, Y. G. Kim, B. H. Lee and S. H. Lee. 2012a. Transgenic expression of MsHsp23 confers enhanced tolerance to abiotic stresses in tall fescue. *Asian-Aust. J. Anim. Sci.* 25:818-823.
- Lee, K. W., J. Y. Cha, K. H. Kim, Y. G. Kim, B. H. Lee and S. H. Lee. 2012b. Overexpression of alfalfa mitochondrial HSP23 in prokaryotic and eukaryotic model systems confers enhanced tolerance to salinity and arsenic stress. *Biotechnol. Lett.* 34:167-174.
- Lee, K. W., K. H. Kim, Y. G. Kim, B. H. Lee and S. H. Lee. 2012c. Identification of MsHsp23 gene using annealing control primer system. *Acta Physiol. Plant.* 34:807-811.
- Yan, J. J., S. Q. Bai, X. Ma, Y. M. Gan and J. B. Zhang. 2007. Genetic diversity of *Elymus sibiricus* and its breeding in China. *Chin. Bull. Bot.* 24:226-231.

Acknowledgement

This work was carried out with the support of “Cooperative Research Program for Agriculture Science & Technology Development (Project No. PJ008599042015)”.