



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Classification of Rangeland Vegetation and Modeling of Vegetation Patterns at the Jabal al Akhdar Mountain, Northern Oman

K. Brinkmann
University of Kassel, Germany

U. Dickhoefer
University of Kassel, Germany

E. Schlecht
University of Kassel, Germany

A. Buerkert
University of Kassel, Germany

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/1-2/18>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Classification of rangeland vegetation and modelling of vegetation patterns at the Jabal al Akhdar mountain , northern Oman


K . Brinkmann¹ , U . Dickhoefer² , E . Schlecht² , A . Buerkert¹

¹Organic Plant Production and Agroecosystems Research in the Tropics and Subtropics , University of Kassel , Germany . E-mail tropcrops@uni-kassel.de , ²Animal Husbandry in the Tropics and Subtropics , University of Kassel and University of Göttingen , Germany

Key words : Hajar mountains , rangeland vegetation , species indicator analysis , two way cluster analysis , discriminant analysis

Introduction The species diversity and the levels of endemism within the northern mountains of Oman are particularly high . Despite some earlier work , data are lacking about the distribution and ecology of different plant species , and also the vegetation response to environmental conditions and land use , particularly grazing . The aim of this study therefore was to describe the species composition , floristic diversity and vegetation patterns of open woodlands at Al Jabal al Akhdar along an altitudinal and a grazing gradient .

Materials and methods The species composition and several environmental variables such as browsing / grazing intensity , sward structure and distance to the settlement were investigated for 62 samples (20×30 m) using a nested plot design . Classification analysis (two-way cluster analysis and indicator species analysis) and ordination methods (DCA) was used to define vegetation types and to identify underlying environmental gradients . The species diversity and the functional diversity were calculated to show the effects of grazing intensity and altitude on biodiversity . A modelling approach based on discriminant analysis and the GIS were used to model the distribution of vegetation types within the study area .



Results The five group stage of the cluster analysis was the most informative , with the maximum number of significant indicators (67) . Floristic and structural differences between groups were mainly due to altitude , followed by topographic location and grazing intensity . Groups were defined on the basis of the dominant species : the *Sideroxylon mascatense*-*Dodonaea viscosa* group on grazed and the *Olea europaea*-*Fingerhuthia africana* group on ungrazed plateau sites at 2 000 m a s l . , the *Ziziphus spina-cristi*-*Nerium oleander* group on wadi sites , and the *Moringa peregrina*-*Pteropryum scoparium* group at 1200 , and the *Acacia gerrardii*-*Leucas inflata* group at 1700 m . The plant species richness followed a unimodal distribution along the altitudinal gradient with the highest number of species in the intermediate altitudinal belt . Altogether , 27% of the species showed a high degree of grazing damage .

Conclusions The main environmental factors , altitude and topographic location , surmounted the grazing effects on plant species richness in the present investigation . Future work should attempt to quantify more clearly the grazing effects with the successional shift in species composition and the regeneration ability .