



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Grassland Diversity Enhances Productivity at Low and High Levels of Nitrogen Addition

Tim Carnus  
*Teagasc Environment Research Centre, Ireland*

J. A. Finn  
*Teagasc Environment Research Centre, Ireland*

Laura Kirwan  
*Teagasc Environment Research Centre, Ireland*

A. Cuddihy  
*Teagasc Environment Research Centre, Ireland*

J. Connolly  
*University College Dublin, Ireland*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/1-2/20>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Grassland diversity enhances productivity at low and high levels of nitrogen addition

T. Carnus<sup>1,3</sup>, J. A. Finn<sup>1</sup>, L. Kirwan<sup>1,2</sup>, A. Cuddihy<sup>1</sup> and J. Connolly<sup>3</sup>

<sup>1</sup>Teagasc Environment Research Centre, Co. Wexford, <sup>2</sup>Eugene Lawler Graduate School of Computing, WIT, Co. Waterford, <sup>3</sup>Environmental and Ecological Modelling Group, School of Mathematical Science, UCD, Dublin 4. E-mail: tim.carnus@teagasc.ie

**Key words:** Biodiversity, Grassland productivity, Nitrogen fertilisation

**Introduction** A central theory of ecology postulates that ecosystem processes in more diverse ecosystems are more stable than in less diverse ones, especially when environmental changes occur. Thus, diversity can provide insurance for some ecosystem function responses, such as biomass production (Yachi & Loreau, 1999). This study quantifies the contribution of species identity and evenness to biomass production in mixed grassland systems under different N addition regimes.

**Material and methods** A split-plot field experiment was established in autumn 2006. The main plot treatment varied plant diversity and the split-plot treatment varied nitrogen (N) application (+155 kgN/ha/yr). All plots received a total of 45 kg/ha/yr N in two applications and were harvested 3 times in 2007. Plant community composition consisted of two grasses (G1: *Lolium perenne* and G2: *Phleum pratense*) and two legumes (L1: *Trifolium pratense* and L2: *Trifolium repens*) sown at proportions defined within a simplex design. Thus, evenness (E), a measure of the distribution of the relative abundance of species in a community, was varied (Kirwan *et al.*, 2007). The basic design consisted of 4 monocultures (E=0), 6 two sp.-mixtures (E=0.67) and 18 four sp.-mixtures dominated in turn by each species (88:4:4:4, E=0.29 and 70:10:10:10, E=0.64), by pairs of species (40:40:10:10, E=0.88) and equally represented at the centroid (25:25:25:25, E=1). The design was repeated at two levels of overall initial abundance (low being 60% of high). Total yield was analysed as a function of species identity and evenness (Kirwan *et al.*, 2007) using linear mixed models to account for the split plot random effects. Likelihood Ratio Tests (LRT) were used to identify significant terms in the model. Analyses were performed in R (version 6.0).


**Results and discussion** Both evenness (LRT  $\chi^2=7.81$ ;  $P=0.005$ ) and N addition (LRT  $\chi^2=47.68$ ;  $P<0.001$ ) had significant and positive effects on total yield (Figures 1 and 2). There was a linear relationship between yield and evenness, which was maximum at the centroid where estimated yield was 1.63 t/ha more than expected from monoculture yields. The interaction between N treatment and evenness (N\*E) was not significant (LRT  $\chi^2=0.91$ ;  $P=0.341$ ). The positive overall effect of N addition depended on the increase in yield of the two grass species (average increase of  $1.14 \pm 0.14$  t/ha in monocultures at high N level; Figure 2).

Unlike other similar studies (Kirwan *et al.*, 2007) transgressive overyielding did not occur; mixtures (E>0) did not consistently yield higher biomass than the best-performing monoculture (Figures 1 and 2). The centroid yield at the low level of N addition was comparable to the yield of the best-performing monoculture at the high N level.


**Conclusions** These results indicate benefits of using agronomic mixtures in managed grasslands under high and low levels of N addition. The positive effect of evenness at both high and low levels suggests that the diversity effect was not solely due to symbiotic N fixation. Other benefits of grassland mixtures will be investigated, and include stability of yield over time and resistance to weed invasion.

### References

- Kirwan L. *et al.* (2007). Evenness drives consistent diversity effects in an intensive grassland system across 28 European sites. *Journal of Ecology*, 95, 530-539.
- Yachi & Loreau (1999). Biodiversity and ecosystem productivity in a fluctuating environment: the insurance hypothesis. *PNAS*, 96, 1463-1468.



**Figure 1** Total yield at different levels of sown evenness. Lines represent high (triangles) and low (circles) levels of N addition.



**Figure 2** Mean (+ sed) yield of species in monoculture and at the centroid (dark grey: low N; light grey: high N).