



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Temporal and Spatial Variation of Arbuscular Mycorrhizal Fungi Spores in Seasonally Salt Stressed Grassland Using *Medicago sativa* for Reclamation in Hexi Corridor, China

Liang Jin
Lanzhou University, China

Chang Wang
Lanzhou University, China

Guoliang Chen
Lanzhou University, China

Yin Zhao
Lanzhou University, China

Xiangwei Sun
Lanzhou University, China

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/1-2/36>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

Liang Jin, Chang Wang, Guoliang Chen, Yin Zhao, Xiangwei Sun, and Xiaojuan Wang

Temporal and spatial variation of arbuscular mycorrhizal fungi spores in seasonally salt stressed grassland using *Medicago sativa* for reclamation in Hexi Corridor, China

Liang Jin, Chang Wang, Guoliang Chen, Yin Zhao, Xiangwei Sun, Xiaojuan Wang*
College of Pastoral Agriculture Science and Technology, Lanzhou University, Gansu Grassland Ecological Research Institute, P.O. Box 61, Lanzhou 730020, China (* Correspondence author's E-mail: xiaojuanwang@lzu.edu.cn)

Key words : arbuscular mycorrhizal fungi, depth, *Medicago sativa*, salinity, season

Introduction Soil salinity is a problem of grave concern because it adversely affects growth and development of plants, especially in arid and semi-arid regions. Arbuscular mycorrhiza (AM) are ubiquitous symbionts between the fungus and host plant. This relationship enhances plant growth through increasing nutrient acquisition and carbon cycling, alleviating environmental stresses, and increasing biomass production. The objective of this study was to investigate the seasonally salt stressed affecting AMF spores of taproot systems plants at the reclamation grassland.

Materials and methods This study was conducted in the Linze Ecological Research Area of Lanzhou University (N39°15', E100°02'), located in Hexi Corridor, Northwest of China. *Medicago sativa* was planted as test materials to reclamation seasonally salt grassland in 2001, 2002, 2003 and 2004, respectively. Rhizosphere soil samples were collected at four different depth as 0-20 cm, 20-50 cm, 50-100 cm and 100-200 cm in April (Shoot), May (Growth), June (Flower), and August (Seed) in 2005. AMF spores were isolated by wet sieving followed by sucrose gradient centrifugation (Daniels & Skipper, 1982). Data were analysed using Univariate Analysis of Variance with SPSS (v13.0).

Results Our results showed that the phenological phase, soil depth, and planted years influence the AMF spores significant (Table 1). (1) The 0-20 cm layer soil has the highest numbers of AMF spores ($9.39 \pm 0.21/g$ dry soil), deeper rhizosphere soil reduced the richness of AMF spores. (2) The longer years that *M. sativa* planted, the more AMF spores were existed. (3) To the phenological phase, the largest number of AMF spores was in June (Flower) ($4.41 \pm 0.21/g$ dry soil), and the lowest was in May (Growth) ($3.20 \pm 0.21/g$ dry soil).

Table 1 Univariate analysis of variance on the effects of growth stage, depth and plant years on the numbers of AMF spores.

Source	Sum of Squares	df	Mean Square	F	Sig
Growth Stage (S)	60.459	3	20.153	5.983	0.001
Depth(D)	3538.322	3	1194.107	354.527	<0.001
Planted Year(Y)	145.085	3	48.362	14.358	<0.001
S * D	169.116	9	18.791	5.579	<0.001
S * Y	73.579	9	8.175	2.427	0.012
D * Y	265.222	9	29.469	8.749	<0.001
S * D * Y	227.446	27	8.424	2.501	<0.001
Error	862.252	256	3.368		

Discussion Mycorrhizal symbiosis is a key component in helping plants cope with adverse environmental condition. In this study, results showed that AMF spores could be used as an indicator to evaluate the degraded ecosystem, and *M. sativa* is a potential materials to reclamation saline soil.

Acknowledgements This study was funded by National Basic Research Program of China (2007CB108902) and Natural Science and Technology Program of Lanzhou University (582402, 582403).