



A Plant-Physiology Approach to a Fire-y Problem

Helen G. Daily
University of Tasmania, Australia

Shaun Lisson
CSIRO, Australia

Kerry L. Bridle
University of Tasmania, Australia

Peter A. Lane
University of Tasmania, Australia

Stuart Anderson
Ministry for Primary Industries, New Zealand

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/22/3-6/6>

The 22nd International Grassland Congress (Revitalising Grasslands to Sustain Our Communities) took place in Sydney, Australia from September 15 through September 19, 2013.

Proceedings Editors: David L. Michalk, Geoffrey D. Millar, Warwick B. Badgery, and Kim M.

Broadfoot

Publisher: New South Wales Department of Primary Industry, Kite St., Orange New South Wales, Australia

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

Helen G. Daily, Shaun Lisson, Kerry L. Bridle, Peter A. Lane, Stuart Anderson, and Ross Corkrey

A plant-physiology approach to a fire-y problem

Helen G Daily^{ABE}, Shaun Lisson^C, Kerry Bridle^A, Peter Lane^A, Stuart Anderson^D and Ross Corkrey^A

^A Tasmanian Institute of Agriculture, University of Tasmania, Pte Bag 98, Hobart, Tas. 7001, Australia.

^B Bushfire Cooperative Research Centre, East Melbourne, Vic. 3002, Australia.

^C CSIRO Ecosystem Sciences, Hobart, Tas. 7001, Australia.

^D MPI Policy Ministry for Primary Industries, Rotorua, 3040, New Zealand.

^E present address: High Resolution Plant Phenomics Centre, CSIRO Plant Industry, GPO Box 1600, Canberra, ACT 2601, Australia.

Contact email: helen.daily@csiro.au

Keywords: Pasture modelling, curing, fire danger rating.

Introduction

As vegetation dies, it dries and becomes more flammable. Fire agencies require accurate and timely assessments of curing (the percentage of dead material in the sward) to model grass fire behaviour and calculate fire danger ratings (Cheney and Sullivan 2008). Visual observation is commonplace and the more objective use of the Levy Rod is recommended, although both have drawbacks (Anderson *et al.* 2011). There is great potential for pasture growth models to provide curing estimates to assist with the management of wild grass fires (Gill *et al.* 2010). This PhD project focused on plant physiological characters to populate models that could be used to predict curing assessments for fire management purposes.

Existing pasture growth models

APSIM (McCown 1996), GrassGro (Moore *et al.* 1997) and the SGS Pasture Model (Johnson *et al.* 2003) were used to simulate curing in a number of pastures (*e.g.* Fig. 1) and a wheat crop in SE South Australia. Curing outputs were similar to estimates derived from the Levy Rod technique except in native pastures. However, no single pasture growth model was suited to all grassland situations.

The models varied in the species simulated, the underlying senescence assumptions, and management flexibility. Use of these models is predicated on accurate inputs and robust validation, which is often problematic. Calibration of curing outputs to management practice is also necessary.

Measurement and modelling of leaf turnover rates

Curing is a fire management term encompassing the senescence and death of plant material, which is not well represented in current plant growth models. Leaf growth rates (leaf appearance, elongation, length, and life span) and leaf senescence rates were measured to capture the whole life cycle to ascertain similarities and differences between four C3 grass species (Fig. 2). This study was unique in that it measured all growth and death stages, including an Australian native, and compared grass species of differing growth habits. Generally, the relationship between leaf rates and leaf position could be represented using non-linear models. This work will be useful to model developers who wish to strengthen the algorithms underpinning particularly the senescence routines in existing pasture growth models. In the interim, a standalone model was developed to incorporate the leaf rates into a prototype curing model.




Figure 1. Comparison of field curing assessments with estimates of curing from GrassGro™ and the SGS Pasture Model for phalaris at Struan, SA, during the 2008-2009 fire season.




Figure 2. Observed (boxplot) and modelled (line) LSR ($10^1 \times \text{mm/gdd}$ ($T_{\text{base}} = 0^\circ\text{C}$)) with leaf position: (a) annual ryegrass; (b) wallaby grass; (c) phalaris; (d) wheat.




Figure 3. Logistic curve based on the Bayesian model for phalaris (line) fitted against visual curing observations from four sub-districts of the Naracoorte Lucindale Council over thermal time (gdd ($T_{\text{base}}=0^\circ\text{C}$) in the 2009-2010 and 2010-2011 fire seasons: ○) Frances; +) Avenue; □) Callendale; ◇) Wrattontully.

Bayesian curing model

The models of individual leaf turnover rates were incorporated into a Bayesian model, to determine the accumulation of green leaf material, as could the progression of curing over time. The Bayesian model for phalaris sufficiently predicted curing in one region (SE of SA) over two fire seasons (Fig. 3).

Conclusion

This project has improved the prediction of curing by applying a plant physiology perspective to what is essentially a plant flammability problem arising from plant senescence and death. This knowledge can be applied to enhance existing tools, or to create new ones from the prototypes developed here.

References

Anderson SAJ, Anderson WR, Hollis JJ, Botha EJ (2011) A

simple method for field-based grassland curing assessment. *International Journal of Wildland Fire* **20**, 804-814.

Cheney P, Sullivan A (2008) 'Grassfires: fuel, weather and fire behaviour.' (CSIRO Publishing: Melbourne, Victoria)

Gill AM, King KJ, Moore AD (2010) Australian grassland fire danger using inputs from the GRAZPLAN grassland simulation model. *International Journal of Wildland Fire* **19**, 338-345.

Johnson IR, Lodge GM, White RE (2003) The Sustainable Grazing Systems Pasture Model: Description, philosophy and application to the SGS national experiment. *Australian Journal of Experimental Agriculture* **43**, 711-728.

McCown RL, Hammer GL, Hargreaves JNG, Holzworth DP, Freebairn DM (1996). APSIM: a novel software system for model development, model testing, and simulation in agricultural systems research. *Agricultural Systems* **50**, 255-271.

Moore AD, Donnelly JR, Freer M (1997) GRAZPLAN: Decision support systems for Australian grazing enterprises. III. Pasture growth and soil moisture submodels and the GrassGro DSS. *Agricultural Systems* **55**, 535-582.