University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress /
VIII International Rangeland Congress

Small-Scale Species Richness and Its Spatial Variation in an Alpine Meadow on the Qinghai-Tibet Plateau

Jun Chen
Shanghai Jiaotong University, China

Huakun Zhou
Chinese Academy of Sciences, China

Masae Shiyomi
Ibaraki University, Japan

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/1-2/3>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Small-scale species richness and its spatial variation in an alpine meadow on the Qinghai-Tibet Plateau

Jun Chen¹, Hua-kun Zhou², Masae Shiyomi³

¹ Agricultural and Biological College, Shanghai Jiaotong University, Shanghai 200240, China. E-mail: chen_jun@hotmail.co.jp. ² Northwest Plateau Institute of Biology, the Chinese Academy of Sciences, Xining 810001, P. R. China.

³ Ibaraki University, Professor Emeritus, Bunkyo 2-1-1, Mito 310-8512, Japan

Key words : grazing, high species richness, *Kobresia* meadow, Qinghai alpine meadow, winter grazing

Introduction We investigated how the high small-scale species richness of an alpine meadow on the Qinghai-Tibet Plateau, China, is maintained. This area is characterized by strong wind and severe cold during long winters. In winter, most livestock is grazed on dead leaves in small pastures near farmers' residences, whereas in the short summer, livestock is grazed in mountainous areas far from farmers' residences.

Study area and methods The number of plant species and the aboveground biomass were surveyed for three adjacent pastures differing in grazing management: a late-winter-grazing pasture grazed moderately from 1 February to 30 April (LWG), an early-winter-grazing pasture grazed lightly from 20 September to late October (EWG), and a whole-year-grazing pasture grazed intensively throughout the entire year (WYG). In each pasture, we harvested the aboveground biomass from 80 or 100 quadrats of 0.01 m² along a transect and classified the contents by species.

Results and discussion We observed 16-20 species per 0.01 m², which is high richness per 0.01 m² worldwide (Table 1). The species richness in the two winter-grazing pastures was higher than that in the whole-year-grazing pasture. The spatial variation in species richness and species composition in the two winter-grazing pastures in which species richness was high was greater than that in the whole-year-grazing pasture in which species richness was lower. Most of the leaves that are preserved on the winter-grazing pastures during summer are blown away by strong wind during winter, and the remaining leaves are completely exhausted in winter by livestock grazing. A pasture with a high richness is accompanied with a high spatial variation in species richness and species composition. There is a high possibility that the characteristics of spatial variation is also caused by the traditional grazing practice in this area.

Table 1 Mean species richness observed in 80 0.01-m² in 2002 and 100 0.01-m² in 2003, variance in richness among quadrats, coefficient of variation (CV) of richness, and the total aboveground richness biomass per 0.01-m² quadrat.

Item	EWG	LWG		WYG	
	2002	2002	2003	2002	2003
Mean species richness (0.01-m ⁻²) ± se	19.36 ± 0.30	19.09 ± 0.33	19.69 ± 0.38	16.11 ± 0.24	15.51 ± 0.30
Evenness for species biomass *	0.770	0.688	0.775	0.727	0.780
Variance for richness (0.01-m ⁻²)	7.22	8.79	14.64	4.51	9.08
Observed CV of richness	0.139	0.155	0.194	0.132	0.194
Theoretical CV of richness **	0.137	0.144	0.146	0.145	0.156
Observed CV/Theoretical CV	1.01	1.08	1.33	0.91	1.24
Biomass (g/0.01m ²) ± se	3.32 ± 0.115	3.10 ± 0.097	3.02 ± 0.081	3.26 ± 0.109	3.14 ± 0.099

Calculated by the formula of $\{\sum p_i(1-p_i)/(\sum p_i)^2\}^{1/2}$, where p_i denotes the occurrence rate per quadrat for species i , and $\sum$ means sum for all species appeared. The theoretical CV-value was calculated under an assumption that each species distribute at random quadrat.

Reference

Chen, J., Yamamura, Y., Hori, Y., Shiyomi, M., Yasuda, T., Zhou, H.K., Li, Y.N., Tang, Y.H., (2007). Small-scale species richness and its spatial variation in an alpine meadow on the Qinghai-Tibet Plateau. *Ecological Research*.