



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Effect of Intercropping Forages on Root Growth of Fruit Tree and Orchard Ecology

Yibin Huang

*Fujian Academy of Agricultural Science, China*

Weibin Pan

*Minxi Vocational and Technical College, China*

Zhaoyang Ying

*Fujian Academy of Agricultural Science, China*

En Chen

*Fujian Academy of Agricultural Science, China*

Boqi Weng

*Fujian Academy of Agricultural Science, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/1-5/7>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Effect of intercropping forages on root growth of fruit tree and orchard ecology

HUANG Yi-bin<sup>1</sup>, PAN Wei-bin<sup>2,1</sup>, YING Zhao-yang<sup>1</sup>, CHEN En<sup>1</sup>, WENG Bo-qi<sup>1</sup>

<sup>1</sup>Eco-agriculture Research Institute, Fujian Academy of Agricultural Science, Fujian Engineering and Technology Research Center for Hilly Prataculture, Fuzhou 350013, Fujian, China. <sup>2</sup>Minxi Vocational and Technical College, Longyan 364021, Fujian, China

**Key words :** forages, *Chamaecrista rotundifolia*, orchard ecology, root system

**Introduction** Intercropping forages in young orchards not only could fully utilize the land resource to solve a conflict between orchard and pasture, but also could improve the microclimate of hilly orchards (Qi et al, 1993). In contrast, some articles reported that planting mulch grasses in orchard could lead to conflicts of competing water and nutrients between grasses and fruit trees (Catzeflis et al, 1997). This paper investigated the root distribution of forage-orchard compound system and effects of intercropping legume forage *Chamaecrista spp.* on the orchard ecology to provide scientific basis for water and fertilizer management of orchard.

**Materials and methods** Experiment 1: The experimental materials were five-year Longan trees and *Chamaecrista rotundifolia* intercropped in Longan tree orchard. Randomized block design was employed. Experiment 2: Four treatments were imposed in a randomized complete block design with 3 replicates. Treatments were: A. Planting *C. rotundifolia* on built contour terraces and Bahia grass and Premier finger grass on the walls of terraces of the orchard; B. No grasses on built contour terraces and the walls of terraces compared with A; C. Planting fruit trees down-slope without building contour terraces, the slope was covered by forages; D. No grasses covered the slope compared with C.

**Results** (1) 88.2% roots of *C. rotundifolia* were concentrated in 0-20 soil layer, however 71.5% roots of Longan tree distributed in the lower soil layer (lower than 0-20cm) (Table 1), thus the overlapped rate of the root system of two different plants was less than 5%. (2) Compared with the weed-cleaning orchard, runoff amount of the red-earth orchard where intercropped *C. rotundifolia* for 3 years decreased from 2482.4t/hm<sup>2</sup> to 61t/hm<sup>2</sup>; the available N, P, K content increased from 47.6, 2.3, 30.4mg/kg to 85.4, 9.0, 95.1mg/kg respectively; and soil porosity and soil moisture increased by 7.1% and 1.2% respectively (Table 2).

**Table 1** Root distribution of 5-year Longan trees (root number/m<sup>2</sup>).

Soil layer(cm)	Distance from the tree base (cm)					Sum	Percentage (%)
	50	100	150	200	250		
0-20	660	550	480	230	140	2060	28.5
20-40	870	800	860	370	180	3080	42.5
40-60	380	550	50	160	100	1240	17.1
60-80	330	120	0	60	70	580	8.0
80-100	130	110	0	0	40	280	3.9

**Table 2** Effect of intercropping *C. rotundifolia* in orchard on soil bulk density and porosity.

Item	Soil layer(cm)	Treatment A	Treatment B
soil bulk density(g/cm <sup>3</sup> )	0-10	0.98	1.05
	10-20	0.97	1.08
Porosity(%)	0-10	25.76	20.1
	10-20	25.70	17.1

**Conclusions** Planting *C. rotundifolia* outside the crown range of Longan trees had little effect on root growth of fruit trees and could reduce the runoff and sediment, raise soil fertility and improve the microenvironment of orchards.

### References

- Qi Zhiping, Tang Shumei, (1996). Tropical forage and eco-agriculture. *Research of Tropical Crops* (3): 30-33.  
 Catzeflis J., (1988). No-tillage of the space between rows in arboriculture compared with grassing down. *Hort Abst* 59(11): 10-13.