



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Character of Canopy Apparent Photosynthesis and Transpiration in *Seriphidium* Semidesert under Different Degradation Gradient

W. H. Lu
Xinjiang Agricultural University, China

J. Z. Zhu
Xinjiang Agricultural University, China

G. L. Jin
Xinjiang Agricultural University, China

D. J. Wang
Xinjiang Agricultural University, China

H. L. Liu
Xinjiang Agricultural University, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/1-5/25>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Character of canopy apparent photosynthesis and transpiration in *Seriphidium* semidesert under different degradation gradient

W-H Lu , J-ZH Zhu , G-L Jin , D-J Wang , H-L Liu

College of Pratacultural Engineering , Xinjiang Agricultural University , Urumqi 830052 China

Key words : Seriphidium semidesert , fence , degradation , CAP , transpiration , daily variation

Introduction *Seriphidium* is generally the dominant native plant in semidesert and is extensively distributed in arid regions , especially on the northern slopes of the Tianshan Mountains . It plays an important role in stockbreeding and ecosystems . Photosynthesis and transpiration , which are two of the primary metabolic processes determining plants growth . But research reports about this semi-desert vegetation CAP and TR are relatively few . The objectives of this study were to determine : 1) what changes under different vegetation CAP and TR in different gradient of degradation in a day ? 2) What relationships between CAP .TR and environmental factors ?

Material and methods The study area is the spring autumn pastures of Sangong village , located on the northern side of the Tianshan Mountains , in Xinjiang , China . The dominant vegetation on the study sites is *Seriphidium* . , the companion species are , *Petrosimonia* and *Ceratocarpus* , but many degraded areas are inhabited by *Petrosimonia* . We divided grassland into three types according to degree of degradation , moderate , serious and exceeding (MD , SD , and ED) and built fence in all types (inside of fence : IF , outside of fence : OF) , with each plot measuring $40 \times 40 m^2$. Photosynthesis was measured using Beijing Siaidi company's CB-1101 photosynthetic detector within an assimilation box measuring $60 \times 60 \times 60 cm^3$ from an aluminum alloy frame and cover of film that allowed light transmittance greater than 85% , within the assimilation box two 12V fans were installed for blending air . Measurements were made from 10 :00am to 18 :00pm , at one hour intervals , three replications were set according fence \times degree of degradation . Photosynthetic and transpiration rate were calculated using the apparatus software package . We inspected the relationships between all the variables by means of correlation analysis . All statistical analyses were performed using the statistical software package SPSS 12.0 .

Results On moderate degraded grassland , with *Seriphidium* as the main vegetation , CAP from 10 :00 to 14 :00 increased slowly . The CAP was $8 \mu mol m^{-2} s^{-1} \sim 9 \mu mol m^{-2} s^{-1}$, and declined rapidly to the minimum from 15 :00~16 :00 , suggesting an obvious " siesta phenomenon" , then rose slowly from 16 :00 to 18 :00 . From 10 :00 to 14 :00 , CAP (IF>OF) , but then both have similar trend in CAP after 14 :00 . When *Petrosimonia* was the main vegetation of severe and exceeding degradation of grassland , both inside and outside of fence CAP changed similarly with an obvious single peak curve with a maximum at 14 :00 . Inside of fence the highest average CAP was with severe degradation , $14.8194 mol m^{-2} s^{-1}$, and the lowest CAP occurred with exceeding degradation outside of fence , only $1.9639 mol m^{-2} s^{-1}$. This result occurred mainly because with the conditions of exceeding degradation , vegetation is damaged and bare soil increases (Figure 1) . TR value of all vegetation types are a single peak curve , related closely to temperature changes . In all vegetation types TR increased significantly at 12 :00 , both its peak appeared in the afternoon at 15 :00 . This result occurred mainly because the transpiration is closely related to change of temperature , which on average TR is the highest on SDIF , followed by EDIF , and the TR changes are smallest on ED OF as the vegetation is very sparse (Figure 2) .

PAR and temperature changes are typical single-peak curves , the changes of temperature lags behind PAR , the peak of PAR appears at noon 14 :00 , but the temperature in the afternoon at 15 :00 . Inside and outside of the fence (Moderate Degradation) CAP decreased and TR increased with increasing PAR . CAP increased significantly with the rise in temperature in serious and exceeding degraded grassland , changes were consistent with temperature and PAR changes , correlation coefficients are 0.95^{**} ($P < 0.01$) and 0.96^{**} ($P < 0.01$) and 0.82^{**} ($P < 0.01$) , 0.34 ($P > 0.05$) , respectively , except with extreme degradation . Due to sparse vegetation CAP did not change significantly with increasing temperature and PAR . Changes of transpiration are closely related to temperature , in the other five cases , the transpiration rate and temperature showed a significant correlation except for extreme degradation , correlation is not significant ; correlation coefficients were 0.80^{**} ($P < 0.01$) , 0.68^* ($P < 0.05$) , 0.87^{**} ($P < 0.01$) , 0.62^* ($P < 0.05$) , and 0.94^{**} ($P < 0.01$) , respectively .




Figure 1 Variation of CAP .




Figure 2 Variation of Transpiration .