



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Effects of Different Factors on the Hypocotyls Protoplast Isolation of Common Sainfoin

Yupeng Luo
Xinjiang Agricultural University, China

Bo Zhang
Xinjiang Agricultural University, China

Aiping Chen
Xinjiang Agricultural University, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/1-5/28>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Effects of different factors on the hypocotyls protoplast isolation of common sainfoin

LUO Yu-peng , ZHANG Bo , CHEN Ai-ping

Xinjiang Key Laboratory of Grassland Resources and Ecology , College of Grassland science of Xinjiang Agricultural University , Urumqi , 830052 , China ,E-mial xjauzb@126 .com

Key words : sainfoin , hypocotyl , protoplast , effect factors , culture

Somatic hybridization which based on protoplast culture has been widely used in forage breeding . In this text , we explore the technology of protoplast isolation , and hope to culture a new pasture cultivars with superior character .

Materials and methods Axenic hypocotyl cultures of common sainfoin . Sainfoin seeds were surface-sterilized with 70% (v/v) ethanol (30s) , then transferred to 0.1% (w/v) mercuric chloride (15min) , and washed 5 times with sterilized water . Seeds were put on half-strength MS basal medium (Murashige & Skoog , 1962) with 0.7% (w/v) agar and 3% (w/v) sucrose . and kept in darkness(25±2°C) for 3-4d days .

They were cut transversely into slice (0.5 mm wide approx .) When the hypocotyls were 2-3cm . The 1-g hypocotyls segments were treated (1 h) in 10 ml CPW salts solution containing 0.7 M mannitol . then transferred into 10 ml filter-sterilized enzyme solution . After 2-10h incubation in the dark (25±2°C) , with gentle shaking (40 rpm) on a rotary shaker , the mixture was passed through a nylon sieve(38.5µm pore sizes) and 15 ml of CPW9M solution was added . The protoplasts were collected by centrifugation (100×g , 5 min) and resuspended in the washing solution . The washing treatment was done twice . Protoplasts , free of debris were carefully removed from the interface of the solutions , and protoplasts were rinsed twice with 15 ml of KM8p medium((Kao and Michayluk , 1975) A small sample of protoplasts in the washing solution was stained with 0.01% (w/v)phenosafranine and yield determined using a haemocytometer .

Results and discussion The principle of enzymolysis is getting viable protoplasts with lower concentration of enzymes and shorter enzymolysis time . Mannitol can adjust osmosis pressure of cell in protoplast isolation . If osmosis pressure too high or low cell membrane will ruptured . pH not only affect the viability of protoplast , but also the activity of enzymolysis . The result showed that the protoplasts with higher yield and quality were obtained by treating the hypocotyls with an enzyme mixture (pH5.8) containing 2% cellulase Onozuka R-10+0.5% Pectinase+0.3% macerozyme R-10 and 0.55mol/L mannitol for 6h .

Table 1 The effects of different enzyme combination on protoplast isolation .

Cellulase Onozuka R-10(g/l)	Pectin-ase (g/l)	Macerozy-me R-10(g/l)	yield of viable protoplast (1×10 ⁶)
10	5	3	0.83
10	5	5	1.18
10	8	3	1.54
10	8	5	1.87
20	5	3	3.20
20	5	5	3.11
20	8	3	2.34
20	8	5	1.96






Figure 1 and Figure 2 The effects of enzymolysis time and mannitol concentration on protoplast isolation .

Figure 3 The effects of pH on rotoplast isolation .

References

- Murashige T & Skoog F (1962) . A revised medium for rapid growth and bioassays with tobacco tissue cultures . [J] *Physiol. Plant* . 56 : 473-497 .
- Kao KN & Michayluk MR (1975) . Nutritional requirements for growth of *Vicia hajastana* cells and protoplasts at a very low population density in liquid media . *Planta* 126 : 105-110 .