



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Grazing Impacts on Natural Steppe Community of Eastern Mongolia

N. Narantuya
Mongolian Academy of Sciences, Mongolia

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/1-5/36>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Dynamics of reproductive allocation of reproductive tillers in *Stipa baicalensis* populations on the Songnen Plains , China

Wang Tie-zhi , Yang Yun-fei

Key Laboratory of Vegetation Ecology , Ministry of Education , Institute of Grassland Science , Northeast Normal University , Renmin Street , 5268 , Changchun 130024 , China . E-mail : yangyf@nenu.edu.cn

Key words : *Stipa baicalensis* , reproductive tiller , biomass , reproductive allocation

Introduction Reproductive allocation (RA) is the biomass ratio of reproductive organs to total biomass in all assimilative resource of a plant individual . Individual is plastic in size due to the difference of the microenvironment . The plasticity follows certain rules and shows the regulation in growth and allocation at the population level . *Stipa baicalensis* , a perennial grass , is usually the constructive or companion species of communities in meadow steppe regions in China (Li , et al . , 2003) . The objective was to find the rules of the relationship of RA and height , biomass of tiller in different growing stages , which provided scientific accumulation to further research of reproductive ecology .

Materials and methods The study was conducted at the Pasture Ecology Research Station of Northeast Normal University , Changling , Jilin province of China (44° 45' N , 123° 31' E) which has temperate sub-humid continental monsoon climate . *S. baicalensis* experimental population was planted in 2005 . Each plot had an area of 2 m × 6 m with 5 replications . Forty reproductive tillers were sampled randomly in all plots at flowering and ripening stages , respectively . Height , tiller and inflorescence biomass were measured . The ratio of inflorescence biomass to tiller biomass was regarded as reproductive allocation . Relationships between biomass and height of the tiller , and RA and the height , the tiller biomass were evaluated using regression analysis .

Results The relationship between biomass and height of the tillers were positively linearly correlated ($p < 0.01$) at flowering and ripening stages , but between RA and both height and biomass of the tiller negative correlation was observed , and only at flowering stage between RA and height of the tiller the correlation was significant ($p < 0.05$) (Figure 1) . These indicated best isometric rule of the biomass and height of reproductive tillers . The rule of negative correlations on RA and height as well as biomass was gradually weakening from flowering to ripening stage . These showed RA of the tillers in different size were all tending to the average value of the population with growing stages .

Discussions The growth , intraspecies and interspecies competitions of plants shared in limited resource , and appropriate combination of growth and reproduction could form relative predominance in evolution . Taller tillers had good light and more nutrition , so they were advantageous in competition . Positive correlation of biomass and height indicated that increasing height was advantageous to the whole tiller , and an increasing ratio of vegetative growth would decrease RA . The nearly accordant RA in ripen embodied the strategies which *S. baicalensis* population advanced fitness through regulating growth and allocation .

References

- Li J . D . , Yang Y . F . (2003) . Analysis of structure types for plant species in *Stipa baicalensis* meadow steppe in the Songnen Plains of China . *Acta Agrestia Sinica* , 11(1) : 15-22 .
 Yang Y . F . , Zhang B . T . (2005) . Module structures and the growth pattern of *Stipa baicalensis* clone in Songnen Plain of China . *Chinese Journal of Applied Ecology* , 16(12) : 2311-2315 .




Figure 1 Observed data and simulated curves on relationship between total biomass (TB) and height(H) of the tiller , RA and both height and total biomass of the tiller at flowering (FS) and ripening (RS) stages .