



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Effects of Cultivation of *Puccinellia tenuiflora* on the Chemical and Physical Characteristics of Alkalinized Soil in Western Areas of Jilin

N. Ishikawa  
*University of Tsukuba, Japan*

M. Kato  
*University of Tsukuba, Japan*

K. Shimizu  
*University of Tsukuba, Japan*

M. Amari  
*National Institute of Livestock and Grassland Science, Japan*

W. Cao  
*Chinese Academy of Agricultural Science, China*

*See next page for additional authors*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/2-1/14>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

**Presenter Information**

N. Ishikawa, M. Kato, K. Shimizu, M. Amari, W. Cao, X. Yu, and A. Xu

## Effects of cultivation of *Puccinellia tenuiflora* on the chemical and physical characteristics of alkalinized soil in western areas of Jilin

N. Ishikawa<sup>1</sup>, M. Kato<sup>1</sup>, K. Shimizu<sup>1</sup>, M. Amari<sup>2</sup>, W. Cao<sup>3</sup>, X. Yu<sup>4</sup> and A. Xu<sup>4</sup>

<sup>1</sup>Graduate School of Life and Environment Science, University of Tsukuba, Tsukuba, Ibaraki, 305-8572, Japan, E-mail: ishikawa@sakura.cc.tsukuba.ac.jp, <sup>2</sup>National Institute of Livestock and Grassland Science, Tsukuba, Ibaraki, 305-0901 Japan, <sup>3</sup>Chinese academy of agricultural science, Beijing, 100081 China, <sup>4</sup>Branch of Animal Science, Jilin Academy of Agricultural Science, Gongzhuling, Jilin, 13610 China

**Key words :** alkalinized soil, alkali-grass, *Puccinellia tenuiflora* chemical and physical characteristics of soil

**Introduction** Alkalinized soils are expanding over the western area in Jilin province of China and have been partially used as grasslands. Alkali-grass (*Puccinellia tenuiflora*), a perennial plant in the *Poaceae* family, grows well in those areas, unlike most grasses which do not grow in soils with very high levels of pH. It has been observed that some grasses become established after several years of cultivation of the alkali grass in these soils. Thus, it is expected that the grass could be used to improve alkalinized soil. Changes in soil characteristics from alkali-grass cultivation, however, were not determined in detail in previous studies. The present study, therefore, was conducted to clarify the effects of alkali-grass cultivation on the chemical components and physical characteristics of alkali soil in Jilin.

**Materials and methods** Soil samples were collected from the alkalinized region around Da-an city of Jilin province. Samples were collected from different depths (0 to 2, 2 to 10, 10 to 20, 20 to 30, 30 to 40 and 40 to 50 cm) from five points in each three types of the alkali areas where alkali-grass had cultivated for three years (Third year) and ten years (Tenth year) or never cultivated (Control), respectively. Electrical conductance (EC mS/cm), pH, minerals (Na, Ca, Mg, K, P, Fe and Cl mg/kg), CO<sub>3</sub><sup>2-</sup> (g/Kg) and HCO<sub>3</sub><sup>-</sup> (g/Kg) of the samples were analyzed after soils were dried and passed through a screen mesh.

**Results and discussion** Mean values of EC, pH (Table 1) and Na contents of shallow layers of soil of Third and Tenth years were lower than those of Control respectively, although those values of middle layer of soil of Third year were higher than those of Tenth year. On the other hand, mean values of EC and Na contents of deep layer of soil of Third year were higher than those of Control. There is significant positive correlation ( $r^2=0.772$ ) between CO<sub>3</sub><sup>2-</sup> content and pH of alkalinized soils. These data suggest that anions in the soils would be neutralized by unknown factor(s) such as organic acids released from roots of alkali-grass and, as a result, sodium concentration of the soils could decrease via transportation of ions.

**Table 1** Characteristics of pH of soils collected from different depth in the alkalinized lands which are cultivated alkali-grass.

Soil type	Depth of soil (cm)					
	0-2	2-10	10-20	20-30	30-40	40-50
Control	11.1	11.4	11.3	11.2	11.3	11.2
Third year	9.9	11.3	11.3	11.3	11.3	11.3
Tenth year	9.9	9.4	9.1	10.1	10.4	10.5

**Conclusion** Based on the results obtained in the present study, it is likely that alkali-grass could improve the alkalinized soil via process of neutralization by unknown factor(s) released from the roots.