



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Effects of Phosphate Fertilizer on Regrowth of *Medicago sativa* after Defoliation

Huixin Jiang  
*Nanjing Agricultural University, China*

Yixin Shen  
*Nanjing Agricultural University, China*

Guiyu Zhai  
*General Station of Animal Husbandry and Veterinary of Shandong Province, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/2-1/16>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Effects of phosphate fertilizer on regrowth of *Medicago sativa* after defoliation

Jiang Hui-xin<sup>1,2</sup>, Shen Yi-xin<sup>1\*</sup>, Zhai Gui-yu<sup>2</sup>

<sup>1</sup> College of Animal Sci. & Tec., Nanjing Agr. Univ., Nanjing 210095; <sup>2</sup> General Station of Animal Husbandry and Veterinary of Shandong Province, Ji'nan 250022. \* Corresponding author, E-mail: yxshen@njau.edu.cn

**Key words:** phosphorus, buds, regrowth, alfalfa, starch

**Introduction** Regrowth after defoliation was related to carbohydrate reserves in resident tissue. Phosphorus advanced regrowth speed and forage yield of alfalfa. A field experiment was conducted to study the effects of phosphate fertilizer on the relationship between regrowth speed and carbohydrate reserves of alfalfa (*Medicago sativa*).

**Materials and methods** The trial was designed randomly with 4 treatments (0, 90, 180, 270 Kg P<sub>2</sub>O<sub>5</sub>/ha) and 3 replicates. The area of plot was 4 m×2 m. Each plot was seeded with 18g alfalfa (var. Sanditi) seeds in 30 cm row space on September 23, 2006. Alfalfa was cut with 5cm stubbles on May 3, 2007. Ten plants from each plot were dug on 0, 2, 5, 9, 13 d after defoliation. The number and dry weight of new buds and shoots, starch content in resident tissue were investigated.

### Results and discussion

#### Regrowth performance




Figure 1 Effects of phosphorus on regrowth.




Figure 2 Starch reserves as influenced by phosphorus.

Buds per plant reached the peak on 5 days after defoliation in all treatments, and mass per plant rapidly increased after 5 days (Figure 1). Both of buds and mass per plant were increased as P application rate increased. The correlation of buds per plant on 5 days and mass per plant on 13 days after defoliation was significant ( $R^2=0.36$ ). It suggested that buds in dormancy were promoted by the application of phosphorus, and then regrowth of alfalfa increased.

**Starch reserves during regrowth** Rapid decrease of starch reserves in root and crown occurred from 0 to 9 days after defoliation (Figure 2). Decreasing rate of starch was higher than that of water soluble carbohydrate. It showed that starch reserves in main root and crown of alfalfa played an important role during early term of regrowth. Starch loss in root during the first 5 days in plot applied phosphate fertilizer rate at 0, 90, 180, 270 Kg P<sub>2</sub>O<sub>5</sub>/ha was 5.1, 8.2, 8.4, 12.9mg/plant, respectively, and the regrowth weight / root starch loss was 33.7%, 36.5%, 55.8%, 52.6%, respectively. The results showed that more starch reserves were utilized and used more efficiently as phosphate application increased.

**Conclusions** Utilization of starch reserves in root and crown of alfalfa during early regrowth period was increased as P application rate increased. Phosphorus could promote the regrowth speed of alfalfa by enhancing the efficiency of root starch reserves for emergence of buds in dormancy after defoliation.

#### References

- Berg, W. K., Cunningham, S. M., Brouder S. M., 2005. Influence of Phosphorus and Potassium on Alfalfa and Yield Components. *Crop Sci.*, 45, 297-304.
- Fu-kuan, Zhai Gui-yu, Shen Yi-xin, 2005. Effect of superphosphate application and rhizobia inoculation on growth and forage quality of *Medicago sativa* in the Yellow River Delta. *Acta Prataculturae Sinica*. 14, 87-93.