



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Sources of Nitrogen in a Napier Grass/Legume Mixture on Smallholder Dairy Farms in Central Kenya

David Miano Mwangi

*Kenya Agricultural Research Institute, Kenya*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/2-1/41>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Sources of nitrogen in a Napier grass/legume mixture on smallholder dairy farms in central Kenya

D. M. Mwangi

Kenya Agricultural Research Institute (KARI) P.O. Box 57811-00200 Nairobi Kenya, E-mail : DMMwangi@kari.org or Mwangi\_mian@yahoo.com

**Key words :** nitrogen , *Pennisetum purpureum* , *Desmodium* , *Macrotyloma* , *Neonotonia*

**Introduction** In Kenya , dairy production is a major source of smallholder farm income , especially in recent years after the collapse of coffee prices . The smallholder dairy systems are characterised by low weight gains in young stock and low milk production (5-6 kg cow<sup>-1</sup> day<sup>-1</sup>) . The low yields are attributed to inadequate year-round feed supply , protein and energy intake . The main source of forage is Napier grass ( *Pennisetum purpureum* ) grown in monoculture . A Napier grass/forage legume mixture can address the constraints to forage yield as the legumes have the ability to fix atmospheric nitrogen (N) through their symbiotic association with rhizobia (Giller , 2001) .

**Materials and methods** The site was in central Kenya (36 .3°E , 0 .30°S) . The soils are humic Nitisols with relatively high inherent fertility . The area receives 1 ,000 mm of rainfall in 2 seasons . The experiment was laid out in a randomised complete block design replicated 5 times with plots measuring 8 x 8 m . Napier grass ( *Pennisetum purpureum* ) cv . Bana was intercropped with one of the following legumes : *Desmodium intortum* cv . Greenleaf (ILRI 104) , *Macrotyloma axillare* cv . Axillare (ILRI 6756) and *Neonotonia wightii* cv . Tinaroo (ILRI 9794) . All the legumes were inoculated with appropriate *Bradyrhizobium* spp . After the first three months establishment period the forage was harvested every 8 weeks and the yield separated into grass and legume components . Biological nitrogen fixation (BNF) was determined using the <sup>15</sup>N natural abundance method .

**Results** The Napier grass/*Desmodium* mixture had the highest legume and total dry matter (DM) yield (  $P > 0 .05$  ) in the three years of the trial , although the grass component yield was slightly lower than that from the mixture with *Macrotyloma* . The mixture with *Neonotonia* had the lowest legume and total DM yield in the same period . Over a period of three years the Napier/*Desmodium* mixture had the highest total nitrogen yield followed by the mixture with *Macrotyloma* (Table 1) . The mixture with *Desmodium* derived the highest amount from biological nitrogen fixation (BNF) while the Napier/*Neonotonia* mixture had the least N from BNF (Table 1) . The proportion of nitrogen derived from BNF was relatively low , the highest being 22% . Even with *Desmodium* which had the highest proportion of N from BNF the mixture still extracted 412 Kg/N ha from the soil (Table 1) .

**Table 1** Nitrogen yield and sources in a Napier grass legume mixture in central Kenya .

Mixture of Napier grass with :	Grass N yield (Kg/ha)	Total N yield (grass+legume) (Kg/ha)	N from BNF (Kg/ha)	Proportion of N from BNF (%)	N from soil (Kg/ha)
Desmodium	265	530	118	22	412
Macrotyloma	353	487	66	14	421
Neonotonia	267	326	32	10	294

**Conclusions** The low proportion of N derived from BNF could be attributed to the high inherent soil fertility at the site . The legume mixtures still extracted a large amount of N from the soil which would have to be replenished but still contributed a large amount of N from BNF .

### Reference

Giller , K . E . (2001) *Nitrogen Fixation in Tropical Cropping Systems* . 2nd Edn (CAB International : Wallingford) .