

## Effects of nitrogen fertilizers on growth and turf quality of Kentucky bluegrass

Yulin Deng<sup>1</sup>, Yan Peng<sup>2</sup>, Yukuan Wang<sup>3</sup>

<sup>1</sup>College of Forestry and Horticulture, Sichuan Agricultural University, Yaan, Sichuan 625000, PRC, E-mail: yulindeng66@126.com, <sup>2</sup>College of Animal Science, Sichuan Agricultural University, Yaan 625000, PRC, <sup>3</sup>Chengdu Institute of Mountain Hazards and Environment, Chinese Academy of Sciences, Chengdu 610041, PRC

**Key words:** *Poa pratensis* cv. Huntsville, nitrogen, fertilization, turf

**Introduction** Interest in turf establishment has boomed in many urban areas of China. Fertilizer application requirements are a key topic in turf management. Various formulations of N-based fertilizers are available for lawn turf. Previous fertilization studies with cool-season turf have used soluble, fast-release N formulations (Moore, R.W., N.E. 1997), or a combination of fast- and slow-release N formulations (Moore, R.W. 1996). Little research has directly compared effects of conventional N fertilizers on introduced Kentucky bluegrass in China.

**Materials and methods** The experimental design was a factorial treatment arrangement with a control (without fertilization) (marked as H<sub>1</sub>, H<sub>2</sub>, H<sub>3</sub>, H<sub>4</sub> and H<sub>5</sub> (CK)) (Table 1) in a randomized complete block (2m × 1m, total 15 blocks) on Kentucky bluegrass (*Poa pratensis* cv. Huntsville). Four conventional nitrogen fertilizers were added (ammonium sulphate, sodium nitrate, ammonium nitrate and urea) under three repetitions. We also fertilized 80 pots (Φ0.23m × 0.20m) with the same treatment regime to determine the biomass yield.

**Results and discussion** (1) Any kind of N fertilization significantly accelerated the foliation velocity and growth rate but without any significant differences among 4 N sources. In the early period after seeding, no significant effect of N fertilizers could be determined on bifurcating ability, but after growth, N fertilization in autumn could greatly improve bifurcation in next spring. Among 4 N sources, ammonium sulphate was the most effective application. Furthermore, turf-formation velocity could be accelerated. Compared with the control, turf on an N fertilized field could be formed 30 days earlier. Admittedly, N fertilization can effectively improve the turf growth and earlier formation. (2) 4 N fertilizers could significantly (p=0.01) deepen the greenness. In spring growing season (see Table 1), 4 N sources made no differences on their effects whereas in autumn growing season (from October to November), Urea application could more significantly increase the leaf chlorophyll content of Kentucky bluegrass. (3) The experiment revealed that 4 N fertilizer applications could significantly increase the biomass of Kentucky bluegrass, but no difference could be testified for effects on aboveground biomass among different fertilizers. (4) Summer withering rate did not reach its peak in the warmest period till late September because of an accumulated high temperature effect. After September 17th, reviving would occur. For different N source applications, any treatment could significantly reduce summer withering rate and ammonium sulphate could be more effective.

**Table 1** Effect of N fertilizers on growth of Kentucky bluegrass.

Treatment	Average foliage		Growth rate (cm/d)		Average bifurcation		leaf chlorophyll content (LCC)				
	Apr 24 <sup>th</sup>	May 14 <sup>th</sup>	Before mowed	After mowed	Sep 14 <sup>th</sup>	Next Apr 14 <sup>th</sup>	May 13 <sup>th</sup>	May 31 <sup>st</sup>	Jun 11 <sup>th</sup>	Oct 1 <sup>st</sup>	Nov 4 <sup>th</sup>
H1	2.87 a	4.43 ab	0.167 a	0.320 a	2.69 a	4.09 a	2.21 A	2.12 A	2.01 A	1.30 AB	1.37 AB
H2	2.80 a	4.02 ab	0.175 a	0.310 a	2.13 a	2.41 bc	2.19 A	2.11 A	1.94 A	1.11 B	1.17 B
H3	2.82 a	4.40 ab	0.180 a	0.360 a	2.50 a	3.77 ab	2.10 A	2.04 A	1.93 A	1.16 AB	1.25 B
H4	2.96 a	4.92 a	0.180 a	0.320 a	2.20 a	3.32 c	2.25 A	2.05 A	1.85 A	1.42 A	1.55 A
CK	2.59 b	3.82 b	0.070 b	0.190 b	2.16 a	1.97 c	1.55 B	1.69 B	1.56 B	0.63 C	0.79 C

**Conclusions** N fertilizers can be effectively applied in turf establishment of Kentucky bluegrass in the sub-tropic area and can improve the foliation, growth of plants and turf formation and quality, and much better in autumn. Moreover, N sources can have different affects. Ammonium sulphate could be more effective for bifurcation and summer tolerance whereas urea for root growth and greenness.

### References

- Mao Tie, Kangfang Xie. (1993). Methodology on turf quality appraisal. [J] *Journal of Grass Science* 8:69-74.  
 Moore, R.W., N.E. Christians, and M.L. Agnew. (1996). Response of three Kentucky bluegrass cultivars to sprayable nitrogen fertilizer programs. *Crop Sci.* 36:1296-1301.