



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Long-Term Phosphorus Fertilization and Perennial Legumes Addition Impacts on a Temperate Natural Grassland: II. Total and Particulate Soil Organic Carbon

Lucía Salvo
Universidad de la Republica, Uruguay

José A. Terra
INIA, Uruguay

Walter Ayala
INIA, Uruguay

Raúl Bermudez
INIA, Uruguay

José Correa
INIA, Uruguay

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/2-2/37>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

Lucía Salvo, José A. Terra, Walter Ayala, Raúl Bermudez, José Correa, Paola Avila, and Jorge Hernández

Long-term phosphorus fertilization and perennial legumes addition impacts on a temperate natural grassland : II . Total and particulate soil organic carbon .

Lucía Salvo¹ , José A . Terra² , Walter Ayala² , Raul Bermudez² , Jose Correa² , Paola Avila² , and Jorge Hernández¹

¹ Departamento de Suelos y Aguas , Fac . de Agronomía , Universidad de la Republica . Garzón 780 , Montevideo , Uruguay , E-mail :lsalvo@fagro .edu .uy , ² Instituto Nacional de Investigación Agropecuaria (INIA) , Ruta 8 Km . 281 , Treinta y Tres , Uruguay .

Key words : Natural grassland , improved pastures , P fertilization , legumes , soil organic carbon .

Introduction Soil organic carbon (SOC) is a critical soil quality indicator and its pools have different functional roles in agroecosystems . Natural grassland improved with perennial legumes and P fertilizers is a rare production system in the world . We evaluated long-term P fertilization and perennial legumes inclusion effects on SOC and particulate organic C (C-POM ; 53-2000 μm) in a natural grassland of Uruguay .

Materials and methods We analyzed a 30 ha experiment in a Typic Argiudol of Uruguay (33° 14' 58" S , 54° 29' 24" W) . A randomized complete block design with five replications was used . Treatments were : natural grassland (NG) , and natural grassland overseeded with *Trifolium repens* L . and *Lotus corniculatus* L . fertilized with 30 (IP₃₀) , or with 60 kg ha⁻¹ yr⁻¹ of P₂O₅ (IP₆₀) during 9 yrs . After the 9th year , soil samples were collected to a depth of 0-5 and 5-15-cm , dried , dispersed and passed through sieves of 2000 , 200 and 53 μm (Cambardella and Elliot , 1992) . Soil remaining on the 200 and the 53 m sieves (coarse and fine C-POM , respectively) and the one passing all sieves (mineral-associated organic matter , C-MAOM) was analyzed for total C using the Mebius method (Nelson and Sommers , 1982) . Orthogonal contrasts were used to make comparisons among treatments , NG vs . IP₃₀ and IP₆₀ average (IP) , and IP₃₀ vs . IP₆₀ . The level of significance for the statistical test was 0 .1 .

Results and discussions Overall , there were no treatments effects on SOC at 0-15-cm depth after 9 yrs (mean : 21 .2 g C kg⁻¹) . However , treatments affected SOC by depth and C pools (Table 1) . On average , IP had 8% higher and 11% lower SOC compared to NG at 0-5-cm and 5-15-cm depth , respectively . On the other hand , IP had 40% higher C-POM (p < 0 .01) and surprisingly 15% lower C-MAOM (p < 0 .01) than NG at 0-15-cm . As expected , a significant C-POM increase of 40% in the 0-5-cm depth was observed in IP relative to NG . In this layer , C-POM represented 46% of total SOC in IP but only 35% of total SOC in NG . Similarly , C-POM , that represented 21% of the total SOC in IP and only 13% in NG at 5-15-cm , was 40% higher in IP than in NG . Unexpected , IP had 9% and 19% lower C-MAOM than NG at 0-5-cm and 5-15-cm depths , respectively . Finally , no significant differences either on SOC or C pools were found between IP₃₀ and IP₆₀ at any depth . The SOC stratification , C-POM increase and C-MAOM drop in IP relative to NG , were probably related with the greater biomass production , lower biomass C-N ratio and the changes in the root system distribution observed in IP . These results are in agreement with Palm et al . (2001) and Metherell (2003) that shown that increased availability of nutrients results in increased quality of the litter input and reduces the recalcitrant pool of SOC .

Table 1 Perennial legumes addition and long-term P fertilization (30 and 60 kg P₂O₅ ha⁻¹ yr⁻¹) impacts on soil organic carbon (SOC) and particulate organic C fractions (C-POM) in a temperate natural grassland of Uruguay after 9 years .

Soil Depth	(0-5-cm)			(5-15-cm)		
	SOC	C-POM (2000-200 μm)	C-POM (200-53 μm)	SOC	C-POM (2000-200 μm)	C-POM (200-53 μm)
	-----g C kg ⁻¹ -----					
Natural Grassland	32 .67b [†]	7 .68b	3 .83b	15 .80a	1 .41b	0 .70b
Improved Pasture 30 kg P ₂ O ₅	36 .02a	11 .44a	5 .15a	13 .96b	1 .75a	1 .33a
Improved Pasture 60 kg P ₂ O ₅	34 .73a	10 .81a	4 .87a	14 .11b	1 .73a	1 .12a

[†] Means followed by the same letter within a column are not significantly different at P ≤ 0 .1 level .

Conclusion The aggregate of data suggest that for temperate NG on undegraded soils , improved pastures with overseeded legumes and P fertilizers may sustain SOC in the long term . However , excessive SOC stratification , basically due to C-POM rise , and C-MAOM drop on IP are issues that may need further attention in the future .

References

- Cambardella , C . A . and Elliot , E . T . (1992) . Particulate soil organic matter changes across a grassland cultivation sequence . *Soil Sci . Soc . Am . J.* 56 :777-783 .
- Nelson , D . W . , Sommers , L . E . , (1982) . Total carbon , organic , and organic matter . In : A . L . Page (Ed) , *Methods of Soil Analysis , Part 2 , Agronomy No . 9* , Am . Soc . Agron . , Madison , WI , pp . 539-579 .
- Metherell , A . K . , (2003) . Management effects on soil organic C in New Zealand pastures . In : *Proc . New Zealand Grassland Assoc.* 65 :253-257 .
- Palm , C . A . , Giller , K . E . , Mafongoya , P . L . , Swift , M . J . (2001) . Management of organic matter in the tropics : translating theory into practice . *Nutrient Cycling in A groecosystem* . 61 :63-75 .