



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Tillage Systems and Biological Nitrogen Fixation of Soybean (*Glycine max*)

Xianzhi Wang
Lanzhou University, China

Yuying Shen
Lanzhou University, China

Tianling Xie
Lanzhou University, China

Murray Unkovich
The University of Adelaide, Australia

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/2-2/46>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Tillage systems and biological nitrogen fixation of soybean (*Glycine max*)

Xianzhi Wang¹, Yuying Shen¹, Tianling Xie¹, Murray Unkovich²

1. College of Pastoral Agriculture Science and Technology, Lanzhou University, Lanzhou 730000, China, E-mail: wangxianzhi@lzu.edu.cn, 2. Soil and Land Systems Group, Earth and Environmental Sciences, The University of Adelaide, Roseworthy SA 5371, Australia.

Key words: tillage system, ¹⁵N natural abundance technology, Loess Plateau

Introduction Legumes play an important role in rotation systems because of their capability in biological nitrogen fixation (BNF), but it is not known if N₂-fixation by legumes is affected by different tillage systems. This paper examines N₂ fixation by soybean (*Glycine max*) in no till, stubble retained and conventional cultivation systems on the Loess Plateau, China.

Material and methods This work was conducted in 2002-2003 on a Heilu soil at Qingyang experimental station on the Loess Plateau, Gansu, China. Four tillage treatments were imposed in a random block design with 4 replications. Treatments were: conventional tillage with no stubble retained (t), conventional tillage but with straw returned to the soil surface after tillage (ts), no-tillage with no straw (nt), and no-tillage with stubble retention (nts). N₂-fixation was assessed using ¹⁵N natural abundance with five Soybean and five Prostrate spurge (*Euphorbia humifusa*) (reference plant) samples/plot. The ¹⁵N natural abundance were analysed by isotope ratio mass spectrometry and the percent of plant N derived from air (% Ndfa) calculated as:

$$\% \text{ Ndfa} = 100 \times (\delta^{15}\text{N}_{\text{ref. plant}} - \delta^{15}\text{N}_{\text{lucerne}}) / (\delta^{15}\text{N}_{\text{ref. plant}} - B)$$

Results In 2002 the nitrogen uptake by soybean were all higher than in 2003 under each treatment, but the significant difference neither in annual variation, nor between treatments. There was no evident difference in % Ndfa between any treatments in 2003 (43.8% ~ 62.4%), but it was much higher than in 2002, and the % Ndfa of soybean was evidently higher in tilled, stubble returned plots (ts) than other treatments. In 2002 the amount of N fixed accounted for 17% ~ 34.4% of the total N uptake by soybean, while in 2003 the quantity of N fixed accounted for 43.3% ~ 66.3% of the N uptake. And the trend of soil NO₃-N (0 ~ 30cm) at sowing was contrary to the % Ndfa and N fixed in 2002 and 2003 (Table 1).

Table 1 Dry matter, dry matter N and N fixed by soybean under different tillage system in 2002 and 2003.

Year	Treatment	Dry matter (kg/ha)	Soil NO ₃ -N (0~30cm) at sowing (kg/ha)	soybean N uptake (kg/ha)	% Ndfa	N fixed by soybean (kg/ha)
2002	t	1299.0	44.1	38.8	17.6	6.6
	ts	1172.3	39.9	33.7	34.3	11.6
	nt	1129.6	45.7	29.1	22.3	6.5
	nts	1179.8	40.2	31.6	19.2	6.1
	LSD _{0.05}	315.2	14.3	11.4	6.0	3.0
2003	t	908.2	21.4	24.7	58.5	14.9
	ts	1024.1	25.3	26.4	62.4	17.5
	nt	825.5	23.8	22.8	54.9	12.9
	nts	936.7	24.7	24.7	43.8	10.7
	LSD _{0.05}	325.7	5.8	9.6	31.0	11.3

Conclusions In the Loess Plateau area, % Ndfa of soybean was increased with a combination of tillage plus straw cover (ts) in 2002 and 2003. There was a significant negative relationship between the quantity of N fixed by soybean and the amount of NO₃-N in the 0~30cm soil profiles. This research suggests that the soil NO₃-N content of 40kg/ha is the critical value above which plant BNF ability is weakened.