



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Effects of K⁺ on Na⁺ Uptake and Accumulation in a Wild Halophyte Grass *Suaeda maritime*

Jinlin Zhang
Lanzhou University, China

Suomin Wang
Lanzhou University, China

Timothy J. Flowers
University of Sussex, UK

Jianyi Cai
Lanzhou University, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/2-3/5>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Effects of K^+ on Na^+ uptake and accumulation in a wild Halophyte grass *Suaeda maritima*

JinLin Zhang¹, SuoMin Wang¹, Timothy J. Flowers², JianYi Cai¹

¹School of Pastoral Agriculture Science and Technology, Key Lab of Grassland Agro-ecosystem, Ministry of Agriculture, Lanzhou University, Lanzhou 730000, China; ²School of Life Sciences, University of Sussex, Falmer, Brighton, Sussex BN1 9QG, UK, E-mail: jlzhang@lzu.edu.cn

Key words: K^+ transporters, $^{22}Na^+$ influx, Na^+ accumulation

Introduction Limiting Na^+ uptake and accumulation in the cytosol is the key measure to improve plant salt tolerance (Flowers, 2004). However, Na^+ uptake pathways in higher plants are not so clear by now. Wang *et al.* (2007) reported that two distinct low-affinity Na^+ uptake pathways exist in *S. maritima*: Pathway 1 might be mediated by a high-affinity K^+ transporter under 25 mM NaCl that is insensitive to TEA^+ or Cs^+ , but sensitive to Ba^{2+} and pathway 2 by an AKT1-type channel under 150 mM NaCl that is sensitive to TEA^+ , Cs^+ , or Ba^{2+} . Here we reported that the turning-point of external salt concentrations for the two pathways and the effects of K^+ on $^{22}Na^+$ influx and accumulation in *S. maritima*.

Materials and methods 17-day-old seedlings were transferred to non- KNO_3 Hoagland solution supplemented with 3 concentrations (0, 10, and 50 mM) of KCl and various concentrations (2.5 to 200 mM) of NaCl for 3 d before being transferred to equivalent solutions labeled with $^{22}Na^+$ or harvested for measurements of influx or concentrations, respectively. Values are means SD ($n=6$ for influx or 8 for concentration test) and bars indicate SD ($P<0.05$, Duncan test).

Results Ba^{2+} inhibited $^{22}Na^+$ influx significantly under various NaCl concentrations, however, TEA^+ showed significant effects in reducing $^{22}Na^+$ influx when the external NaCl concentration was above 100 mM (Figure 1). 10 mM KCl had no significant effect on Na^+ influx under various NaCl concentrations, but 50 mM KCl blocked $^{22}Na^+$ influx significantly when NaCl concentration was above 100 mM (Figure 2). 10 and 50 mM KCl both reduced NaCl concentration in roots under various NaCl concentrations (Figure 3), however, NaCl concentrations in shoots were reduced by 50 mM KCl when external NaCl concentration was below 75 mM and by 10 mM KCl when below 25 mM (Figure 4).




Figure 1 Effects of TEA^+ and Ba^{2+} on $^{22}Na^+$ influx.




Figure 2 Effects of K^+ on $^{22}Na^+$ influx under various NaCl concentrations.




Figure 3 Effects of K^+ on Na^+ accumulation in roots.




Figure 4 Effects of K^+ on Na^+ accumulation in shoots.

Conclusions The turning-point of external salt concentrations for the two pathways was between 75 and 100 mM. K^+ (10 or 50 mM) had no effect on $^{22}Na^+$ influx at concentrations below 75 mM NaCl, but $^{22}Na^+$ influx was inhibited by 50 mM K^+ when the external concentration of NaCl was above 75 mM. However, the affecting mode of K^+ on NaCl concentration in both roots and shoots with increasing external concentration was different from that on $^{22}Na^+$ influx.

Reference

Flowers T. J. (2004) Improving crop salt tolerance. *Journal of Experimental Botany* 55: 307-319.
Wang S. M., Zhang J. L., Flowers T. J. (2007) Low-Affinity Na^+ Uptake in the Halophyte *Suaeda maritima*. *Plant physiology*, 145: 559-571.