

Correlations between soil properties and plant growth for special synthetic soils added with five components used in high-and-cut rock slopes

Gui-juan Gao¹, Rui-Hong Han²

¹Dept. of Biology, Guangdong Education Institute, Guangzhou Guangdong 510303, China

²College of Life Science, South China Agriculture University, Guangzhou, Guangdong Province 510642, China

Key words : rock slope, outside soil spray seeding, orthogonal array designs, synthetic soils, plant, restoration

Introduction It is well known that the outside soil spray seeding (OSSS) is most effective among the various techniques used for re-vegetation of bare rock slopes because of its high mechanization, high efficiency and fast restoration (Albaladejo-Montoro, 2000). However, study of Esther (2004) showed that though OSSS behaved well on the low and gentle rock slopes whose gradients are less than 45°, it was not effective on high-and-cut rock slopes. Previous studies on OSSS were mainly focused on the improvement of engineering techniques (Fowler and Maddox, 1974; Carr and Ballard, 1980). However, the physico-chemical and biological properties of the soils after added different components used for OSSS remain less known. Therefore, We designed a few of synthetic soils with five variables by using orthogonal array method to study the properties of the special synthetic compound soils.

Materials and methods Twenty-five treatments were assigned to tray-planted (40 cm × 30 cm × 15 cm) experiments by using an orthogonal array designs OAD₂₅ (5⁵) matrix with five levels for each factor and five replicates for each treatment. Italian ryegrass was selected as the tray-planted species to assess the status of plant growth. Soil samples for each treatment were collected by using soil core rings (30 mm in diameter and 50 mm in length). In the experiment, plant biomass, soil pH, organic carbon, total nitrogen and phosphorus and soil invertase activities were used as response functions, respectively. Plant biomass was dried at 65 °C to constant weight and expressed on a weight basis. Above ground biomass was measured with the clipping method at plant maturity and below ground biomass was measured with the washing method. PH was measured by Glass calomel Electrode (Smith and Doran, 1996). Organic carbon was measured by the methods of Mebius (1960). Total nitrogen and phosphorus (Bremner and Mulvaney, 1982). Enzyme activities was analyzed using the method of Schinner and Mersi (1990).

Results The Pearson *s* correlation coefficient was used to quantify the strength of relationships existing among all the indices. The *r*-matrix showed that most indices exhibited high and significant relationships between plant growth, soil physico-chemical and biological properties at all concentrations of the five application factors. The plant biomass and showed highly significant and positive correlations with invertase as well as negative correlations with soil pH, total nitrogen and total phosphorous. These results indicated that the high pH and the over-high soil available fertilizers have hurt the plant growth. Considering the correlations between soil physico-chemical and biological properties, soil pH was negatively and significantly correlated to invertase activity, whereas organic carbon showed positively and significantly correlation with invertase activity, which suggested that the effects of soil psyco-chemical properties on soil enzyme activities were significant.

Discussion From the results, we can see that the additive of five amendments significantly affected plant growth and soil physico-chemical and biological properties.

References

- Albaladejo-Montoro, J., Alvarez-Rogel, J., Qurerjeta, J., Díaz, E., Castillo, V., 2000. Three hydro-seeding revegetation techniques for soil erosion control on anthropic steep slopes. *Land Degrad. Dev.* 11, 315-325.
- Esther, B., Patricio, G.F., 2004. Factors controlling vegetation establishment and water erosion on motorway slopes in Valencia, Spain. *Restor. Ecol.* 12, 168-174.
- Fowler, D.K., Maddox, J.B., 1974. Habitat Improvement Along Reservoir Inundation Zones by Barge Hydroseeding. *J. Soil Water Conserv.* 29, 263-265.
- Smith, J.L., Doran, J.W., 1996. Measurement and use of pH and electrical conductivity for soil quality analysis. In: Doran, J.W., Jones, A.J. (Eds.), *Method. Assessing Soil Qual. WI.*, pp. 169-185.