



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Role of Grazing Cattle on Seed Dispersal of Plants in a Hill Pasture 2. Effects of Ruminal Digestion on Seed Germinability of Five Plant Species

M. Obara
Tohoku University, Japan

Shin-ichiro Ogura
Tohoku University, Japan

T. Shishido
Tohoku University, Japan

K. Sugawara
Tohoku University, Japan

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China in 2008.

Edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Role of grazing cattle on seed dispersal of plants in a hill pasture 2 . effects of ruminal digestion on seed germinability of five plant species

M . Obara , S . Ogura , T . Shishido and K . Sugawara

Graduate School of Agricultural Science , Tohoku University , Kawatabi , Naruko-onsen , Osaki , 989-6711 Japan . E-mail : s-ogura@bios.tohoku.ac.jp .

Key words : abomasum , digestion , microbial attachment , rumen , seed dispersal

Introduction Plant seeds , after ingested by grazing herbivores , are exposed to gastro-intestinal digestion . While there have been reports on the change of germination rate of plant seeds by passage in the alimentary tract (e .g . , Ocumpaugh and Swakon , 1993) , there is still scarce information on the effects of ruminal (microbial) and abomasum (enzymatic) digestion on seed germinability of herbaceous plants . The aim of this study was to examine how the ruminal and abomasal digestion affect seed germinability of 5 major plant species common to Japanese hill pastures .

Materials and methods A hundred mature seeds of *Dactylis glomerata* , *Trifolium repens* , *Zoysia japonica* , *Anthoxanthum odoratum* and *Carex albata* were exposed to a two-stage *in vitro* digestion technique (Tilley and Terry , 1963) . To simulate ruminal digestion (step I) , the seeds were incubated with grass hay in dilute ruminal inoculum (20% of strained ruminal fluid collected from a single dairy cattle fed grass silage and 80% of mineral buffer solution) under anaerobic condition for 24 or 48 hrs . To simulate post-ruminal digestion (step II) seeds were incubated for 8 hrs in 0.2% Pepsin-HCl solution . After these treatments , germination rate of control and treated seeds were measured according to methods described in the previous study (Obara *et al.* , 2008) . Seed surface characteristics were observed by using a scanning electronic microscopy (SEM) .

Results and discussion Germination rate of *D. glomerata* , *Z. japonica* , *A. odoratum* and *T. repens* significantly decreased by the combination of step I and II , and 48 hrs incubation at step I ($P < 0.05$) (Figure 1) . In contrast , significantly greater germination rates were observed for longer incubation times in *C. albata* seeds . There may be some factors which promote the germinability of *C. albata* seeds in the rumen and abomasums , such as acidic and enzymatic seed sacification . From SEM we observed few ruminal bacteria attached to the surface of *C. albata* seeds whereas attachments to the surface of the other 4 plant seeds were numerous . This indicates that seed surface of *C. albata* is resistant to microbial digestion while post-ruminal digestion may contribute to increasing germinability .

Conclusions Our results indicate that ruminal and abomasal retention promotes germinability of *C. albata* seeds . Seed surface characteristics that inhibit microbial attachment in the rumen probably have a important role promoting ruminant dispersal of *C. albata* seeds via the gastro-intestinal tract .

References

- Role of grazing cattle on seed dispersal of plants in a hill pasture 1 . Effect of sward structure on seed ingestion by cattle in 3 monocotyledons . *Proc . IGC/IRC Congress , Huhhot , China* . (in press)
- Ocumpaugh , W . R . , Swakon , D . H . D . , 1993 . Simulating grass seed passage through the digestive system of cattle : a laboratory technique . *Crop Science* , 33 : 1084-1090 .
- Tilley , J . M . A . , Terry , R . A . , 1963 . A two-stage technique for the *in vitro* digestion of forage crops . *J Brit Grassl Soc* , 18 : 104-111 .




Figure 1 Germination rate of the seed from 5 plant species post *in vitro* ruminal (step I) and pepsin/HCl (step II) digestion .

Mean values with different letters differ ($P < 0.05$, Tukey's test) .