



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## **Vegetation Response to Grazing Management in a Mediterranean Grassland: A Long-Term Synthesis**

Marcelo Sternberg  
*Tel Aviv University, Israel*

Zalmen Henkin  
*Agricultural Research Organization, Israel*

Avi Perevolotsky  
*Agricultural Research Organization, Israel*

Eugene D. Ungar  
*Agricultural Research Organization, Israel*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/3-1/45>

**The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.**

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## **Vegetation response to grazing management in a Mediterranean grassland : a long-term synthesis**

Marcelo Sternberg<sup>1\*</sup>, Zalmen Henking<sup>2</sup>, Avi Perevolotsky<sup>2</sup> and Eugene D. Ungar<sup>2</sup>

<sup>1</sup>Department of Plant Sciences, Weizmann Institute of Science, Tel Aviv University, 69978, Israel. <sup>2</sup>Department of Natural Resources, ARO, The Volcani Center, PO Box 6, Bet Dagan 50250 Israel. \* Author for correspondence, fax: +972 3 6405877. E-mail: MarceloS@tae.ariel.ac.il

**Key words :** grazing, long-term, Mediterranean grassland, plant functional groups

**Introduction** A long-term synthesis (13 years) on the effects of cattle grazing management on the structure and composition of a Mediterranean grassland in north-eastern Israel is presented. This study provides new insights on the response of the vegetation to rainfall and grazing management.

**Methods** The relationships among plant functional groups was studied in the context of the effects of grazing pressure of the most recent and former grazing seasons, as well as on the rainfall amounts of the most recent and previous rainfall seasons. Treatments included manipulations of stocking rates (moderate, heavy and very heavy) and of grazing regimes (continuous vs. seasonal), in a factorial design.

**Results** Inter-seasonal rainfall variation was a dominant factor in the expression of different grazing treatments on the structure of the plant community. Species diversity was significantly affected by grazing treatments and their effects were stronger in years with dry springs. Grazing effects were stronger on tall annual grasses and annual legumes in wet rainfall years. In dry springs and years, an increase in plant cover was noted in crucifers and thistles with increasing grazing intensity. A reduction in cover of tall grasses was correlated with an increase in cover of less palatable groups such as annual and perennial thistles as well of prostrate and shorter groups such as annual legumes and short annual grasses. Cover of functional groups composed by hemicryptophytic species were less variable (lower CV), in response to grazing treatments compared to functional groups with annual species.

**Conclusions** Increasing grazing intensity produced a shift in dominance of less palatable functional groups and was rainfall dependant. However, persistency of tall grasses and more palatable species support the idea that Mediterranean grasslands are highly resilient. This long-term study shows that the community is rather stable in spite important variation in grazing pressures and rainfall conditions. Grazing pressure could be increased compared to current management pressures, however, the effects of rainfall conditions should be included in the managerial protocols to prevent a reduction in forage quality of the grassland.