



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII  
International Rangeland Congress

---

## Dynamics of Module Structures in *Puccinellina tenuiflora* Clones in Alkalinized Meadow in the Songnen Plains, China

Xiaodong Zhuang  
*Northeast Normal University, China*

Yunfei Yang  
*Northeast Normal University, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/4-1/42>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Dynamics of module structures in *Puccinellina tenuiflora* clones in alkalized meadow in the Songnen Plains , China

Zhuang Xiao-dong , Yang Yun-fei

Key Laboratory of Vegetation Ecology , Ministry of Education , Institute of Grassland Science , Northeast Normal University , Renmin Street , 5268 , Changchun 130024 , China , E-mail : yangyf@nenu.edu.cn

**Key words :** *Puccinellina tenuiflora* , clone plant , module , vegetative tiller , reproductive tiller , alkalized meadow , structure dynamics

**Introduction** *Puccinellina tenuiflora* , a perennial tuft grass (Puccinellia) , is broadly distributed in alkali meadows on the Songnen Plains , tolerates salt and alkali stress , and forms large-area monodominant communities in alkali-patches that have low levels of surface soil (Yang et al , 1995) . Previous studies have characterized vegetative propagation and sexual reproduction , the effect of different methods of utilization on the dormancy , and phenotypic plasticity and prolonged reproductive growth (Yang et al , 1995 ; Zhang et al , 2006 ; Sun et al , 2007) . The objective of this study was to characterize the relationships between vegetative and , reproductive tiller production and their biomass at heading and ripening stages of *P. tenuiflora* clones . These relationships may improve our understanding of *P. tenuiflora* population ecology and provided scientific guidance for grassland management in this region .

**Materials and methods** The research was conducted in natural alkali meadows , located at the Pasture Ecology Research Station of Northeast Normal University , Changling , Jilin province of China (44°45'N , 123°31'E) which has a temperate , half-humid , continental-monsoon climate . Sample plots were located in mono-communities of *P. tenuiflora* . Whole clones were sampled at heading and ripening stages of *P. tenuiflora* . Thirty clones were taken from random locations at each growth stage . The tuft diameters were measured , and the number and biomass of total tillers , vegetative tillers and reproductive tillers were counted and weighted , respectively . The number and biomass of each module were analyzed and tested by one-way ANOVA .

**Results** The tuft diameters of *P. tenuiflora* clones were  $11.0 \pm 1.9$  cm and  $13.6 \pm 2.2$  cm at heading and ripening stages , respectively . The numbers and percents of vegetative and reproductive tillers were not significant ( $p > 0.05$ ) at two stages (Table 1) , indicating the quantitative structures of clones didn't change with growth stages . The biomass of each module was significantly higher at ripening stage than that at heading stage ( $p < 0.05$ ) , and the biomass of vegetative and reproductive tillers at ripening stages were 1.56 and 1.45 times compared with that at heading stage . The differences in percent of the biomass in vegetative and reproductive tillers was not significant ( $p > 0.05$ ) . This indicated that the biomass structures of *P. tenuiflora* clones didn't change during different growth stages .

**Table 1** Quantitative characters of the modules of *P. tenuiflora* clones in different growth stages .

Quantitative characters	Growth period	Vegetative tiller		Reproductive tiller		Total	
		Mean±SD	Percent (%)	Mean±SD	Percent (%)	Mean±SD	Percent (%)
Tillers (clone <sup>-1</sup> )	Heading	33.6±15.0a	32.3±79.0a	72.4±29.1a	67.7±0.1a	106.0±33.9a	100
	Ripening	32±19.0a	30.3±0.1a	70.7±26.3a	69.7±0.1a	102.7±35.2a	100
Biomass (g·clone <sup>-1</sup> )	Heading	1.7±0.9a	15.4±0.1a	8.7±4.7a	84.6±0.1a	10.5±4.0a	100
	Ripening	2.6±1.8b	17.5±0.1a	12.6±4.26b	82.5±0.1a	15.2±4.6b	100

**Conclusions** The reproductive tillers were dominant in *P. tenuiflora* clones , and the dominance of biomass was much more obvious than that of tiller numbers . From heading to ripening stage , both the number and biomass of *P. tenuiflora* clonal modules were relatively stable . The clones could regulate the vegetative propagation , sexual reproduction , production and allocation of nutrition substance as plant populations .

### References

- Sun J . , Yang Y . F . (2007) . Phenotypic plasticity and its regulation of tillers prolonged reproductive growth of *Puccinellia tenuiflora* population on alkalized meadow in Songnen Plains of China . *Chinese Journal of Ecology* , 18(4) : 771-776 .  
 Yang Y . F . , Zhu L . , Li J . D . (1995) . Quantitative characters of vegetative propagation and sexual reproduction of *Puccinellia tenuiflora* population on alkalized meadow of Songnen Plain , China . *Chinese Journal of Applied Ecology* , 6(2) : 166-171 .  
 Zhang L . H . , Zhao J . M . , Yang Y . F . (2006) . Influences of the dormancy modules of *Puccinellia tenuiflora* population under using for grazing and cutting in alkaline meadow . *Pratacultural Science* , 23(4) : 8-11 .