



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Biodiversity, Vegetation Measurements, and Rehabilitation of Foothills of Khanasser Valley (Southeast of Aleppo)

A. Khatib Salkini

International Center for Agricultural Research in the Dry Areas, Syria

M. Audat

Atomic Energy Commission, Syria

J. Tiedeman

International Center for Agricultural Research in the Dry Areas, Syria

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/4-1/43>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Biodiversity , vegetation measurements , and rehabilitation of foothills of Khanasser Valley (Southeast of Aleppo)

A . Khatib Salkini¹ , M . Audat² , J . Tiedeman¹

¹ International Center for Agricultural Research in the Dry Areas (ICARDA) , Aleppo , Syria ; ² Atomic Energy Commission (AEC) , Damascus , Syria , E-mail : A_khatib@cgiar.org

Key words : biodiversity , vegetation measurements , rehabilitation

Introduction The Khanasser Valley is located in northwestern Syria . The current biodiversity of this area is degraded and unpalatable species dominate this area (*Peganum harmala* , *Noaea mucronata*) . Two sites were selected in the foothills of Khanasser Valley to study the biodiversity , herbaceous and shrub biomass , species frequency , species density ; species cover% , and economic uses of native plants . At each site were three treatments : 1) open grazing , 2) fenced plot , and 3) improved fenced plot . The main objective of the study was to evaluate rehabilitation of degraded areas by planting adapted and palatable species of fodder shrubs and perennial grasses and legumes such as : *Poa bulbosa* , *Dactylis glomerata* , *Medicago radiata* , *Salsola vermiculata* , and *Atriplex halimu* . The study was conducted during 1999-2004 .

Material and methods The study involved determining : 1) plant diversity , 2) vegetation characteristics (vegetation cover% , biomass of herbaceous and shrubs , species composition) , 3) plant community characteristics , 4) plant uses , and 5) selection of adapted and palatable species for improving sites .

Results A total of 120 species were observed in the fenced plot and the use of these species (palatable for sheep , medicinal plant , food , fuel , prevent erosion) were recorded . Only 50 species were found in the open grazing plot , and most of these species were unpalatable and poisonous .

Table 1 Multiple uses of major native species .

| Species | Forage | Food | Improving soil fertility | Medicinal | Prevent erosion |
|-------------------------|--------|------|--------------------------|-----------|-----------------|
| <i>Medicago radiata</i> | ✓ | | ✓ | | |
| <i>Hordeum murinum</i> | ✓ | | | | |
| <i>Teucrium polium</i> | ✓ | | | ✓ | ✓ |
| <i>Capparis spinosa</i> | | ✓ | | ✓ | ✓ |
| <i>Peganum harmala</i> | | | | ✓ | ✓ |

Herbaceous biomass in the fenced plot was 485 kg/ha and in the open grazing plot was 100 kg/ha . Shrub biomass in the improved plot was 200 kg/ha , in the natural fenced plot was 40 kg/ha , while in the open grazing area was 7 kg/ha . The most common plant community contained *Hordeum murinum* and *Teucrium polium* . The highest species composition% was observed for *Hordeum murinum* with 42% and *Noaea mucronata* with 27% .




Figure 1 Biomass (t/ha) for herbs , shrubs at two sites .




Figure 2 Plant , stone , soil , bare cover% for two sites .

The best adapted and palatable species on the improved site included : trees (*Pistacia atlantica*) , shrubs (*Atriplex halimus* , *Artemisia herba-alba* , *Haloxylon aphyllum* , *Salsola vermiculata*) , perennial grasses (*Oryzopsis miliacea* , *Phalaris tuberosa* , *Dactylis glomerata*) , and annual legumes (*Medicago radiata* , *Trifolium tomentosum* , *Astragalus asterias* , *Trigonella monspeliaca*) .

Reference

Given , D . R . , (1994) . Principles and Practice of Plant Conservation , Lincoln University , New Zealand . London : Chapman and Hall .