



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Developing a Decision Support Tool for the Steppe Zone of Ghom-Iran, by Implementing a State and Transition Model within a Bayesian Belief Network

H. Bashari
University of Tehran, Iran

C. S. Smith
University of Tehran, Iran

O. J. H. Bosch
University of Tehran, Iran

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/5-1/13>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Developing a decision support tool for the steppe zone of Ghom-Iran , by implementing a state and transition model within a bayesian belief network

H . Bashari , C .S . Smith , O .J .H . Bosch

Faculty of Natural Resources , the University of Tehran , Karaj , Iran , E-mail : H _ bashari2001@yahoo .com

Key words : qualitative knowledge , state and transition model , Bayesian Belief Network , Iran

Introduction Although State and Transition Models (STM) provide a description of rangeland dynamics , the typical descriptive flowcharts and associated catalogue of states and transitions lack practical application . They also handle uncertainty associated with transitions poorly . It is therefore clear that a mechanism is needed to convert these models into predictive models that can accommodate uncertainty associated with the nature of transitions . Bayesian Belief Networks (BBNs) (Jensen 1996) can be used to overcome this problem . The approach is based on BBNs , which are cause and effect models that relate variables through the use of conditional probabilities . This allows for uncertainty to be incorporated into the models . This paper describes the development of a rangeland decision support tool by combining BBN with STM for the Steppe zone of Ghom , Iran .

Materials and methods The STM was the starting point for model development . An influence diagram was built to show the possible transitions and the factors influencing each transition . Next , the influence diagram was populated with probabilities to produce a predictive model , and finally the behaviour of the model was tested using scenario and sensitivity analysis .

Results The STM consists of 7 vegetation states and 15 transitions . The sensitivity analysis revealed that grazing impact and growing condition were the two most important drivers of almost all but two of the transitions . Grazing impact represents the management influence on transition and growth condition represents the environmental influence on transitions . This result is supported by other studies in Iran , which suggest that frequent droughts coupled with mismanagement (e .g . overgrazing) combine to produce rapid land degradation (Nemati 1986) . This result , however , does not match the beliefs of governors or livestock managers . Most governors believe that grazing is the dominant factor responsible for rangeland degradation , while livestock managers believe that it is drought and growing conditions .




Figure 1 This figure shows that within a five-year time frame , the transition to "Unpalatable forbs & shrubs and annuals" is most likely if grazing pressure is high and growing condition is poor . This is most likely where stocking rate is high and soil water availability is low . The numbers are probabilities . Under this scenario , for example , there is a 61 .3% chance that rainfall is less than average .

Conclusions Combining STMs and BBN provides a novel means of modeling rangeland dynamics in that it (a) accommodates uncertainty in rangeland dynamics in a straightforward way , (b) captures experiential knowledge of outcomes of rangeland management scenarios , (c) provided a graphical and transparent modeling environment that can facilitate communication about cause and effect , and (d) allows for the consequences of rangeland management decisions to be predicted . It also identifies the management scenarios that are most likely needed to prevent decline in rangeland condition .

References

Jensen , F .V . , (1996) . An introduction to Bayesian Networks , University College London Press .
 Nemati , N . , (1986) . Pasture improvement and management in arid zones of Iran . *Journal of Arid Environments* , 11(1) :27-35 .