



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Growth and P Uptake of *Dactylis glomerata* L . and *Anthoxanthum odoratum* L . Response to Mycorrhizal Inoculation in Acid Condition

Wu En
Inner Mongolia Agricultural University, China

G. R. Xin
Sun Yat-sen University, China

K. Sugawara
Tohoku University, Japan

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/3-2/41>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Growth and P uptake of *Dactylis glomerata* L. and *Anthoxanthum odoratum* L. response to mycorrhizal inoculation in acid condition

Wu En^{1,3}, G.R. Xin^{2,3}, K. Sugawara³

¹College of Ecol. and Env. Sci., Inner Mongolia Agric. Univ., Huhhot, Inner Mongolia 010018 P.R. of China; ²College of Life Sci., Sun Yat-sen Univ. Guangzhou, Guang dong 510275 P.R. of China; ³College of Agric. Sci., Tohoku Univ., Sendai 981-0914 P.R. of Japan. ¹E-mail: wuen2004@163.com

Key words : acid condition, *Anthoxanthum odoratum* L., mycorrhizal inoculation, *Dactylis glomerata* L., response

Introduction We found the AMF colonization of *A. odoratum* not correlated with soil pH, and coverage of *A. odoratum* strongly correlated with AMF colonization in our field study (Wu et al. 2006). In addition, we also found the mycorrhizal colonization, total spore number in rhizosphere of *A. odoratum* prominently higher than that of the *D. glomerata*, respectively. The spore density of *A. odoratum* and *D. glomerata* were of 15 and 4 spores per 10g fresh soil, respectively. Four different morphological types of spores occurred in rhizosphere of *A. odoratum* and only one morphological type of spore occurred in rhizosphere of *D. glomerata*. These results suggest that the *A. odoratum* invade and extend in severity acidic pasture origin of this specificity on arbuscular mycorrhizal symbiosis in acidic soil other than *D. glomerata*. In present study, we explored the effect of soil pH in mycorrhizal colonization, P uptake and growth of both plants by glasshouse experiment.

Materials and methods Plant materials: *D. glomerata*, is grass; *A. odoratum*, is weed. Treatment and soil modulate: all experiment include 2 (soil sterilized and non sterilized) x 4 (pH=3, 4 (control), 5, 6) = 8 treatments. Cultivating soil collected from an acidic artificial pasture (pH=4), soil sterilized by benomyl (Fitter and Nichols 1988), soil pH modulated by NaOH. Planting and harvest: 4 seedlings of *D. glomerata* and *A. odoratum* mixed planting to per pot respectively at 2 Jun 2003, all treatment harvested at 2 September 2003. Measures: The roots were stained with trypan blue / lactic acid and the mycorrhizal colonization (percentage of root infected by arbuscular mycorrhizal fungi) was calculated using the line-intersect method (Giovannetti and Mosse 1980). Shoot biomass and P content measured after drying at 70°C with 48h. Experiment conducted in farm of college of Agricultural Science of Tohoku University, Osaki, Miyagi prefecture, Japan (38°45'N, 140°45'E).

Results Significantly positive correlation occurred in soil pH between Mycorrhizal colonization of *D. glomerata*, but the *A. odoratum* was not yet. A significantly positive correlation occurred in mycorrhizal colonization between shoot dry mass of both plants. Also a significantly positive correlation occurred in mycorrhizal colonization between shoot P content of both plants (Table 1).

Table 1 Regression equations of some parameters of *D. glomerata* and *A. odoratum*.

Plants	regression equations		
<i>D. glomerata</i>	$y_1 = 10.07x_1 - 24.70$, $r = 0.894$, $p < 0.0001$	$y_2 = 0.01x_2 + 0.23$, $r = 0.867$, $p = 0.0003$	$y_3 = 0.04x_2 + 2.32$, $r = 0.748$, $p = 0.005$
<i>A. odoratum</i>	$y_1 = 1.80x_1 + 20.05$, $r = 0.243$, $p = 0.446$	$y_2 = 0.01x_2 + 0.21$, $r = 0.778$, $p = 0.0029$	$y_3 = 0.07x_3 + 1.61$, $r = 0.797$, $p = 0.002$

y_1 Mycorrhizal colonization, x_1 soil pH, y_2 shoot dry mass, x_2 Mycorrhizal colonization, y_3 shoot P content, x_3 Mycorrhizal colonization.

Conclusions Based on these results, we concluded that the *A. odoratum* have a specific mycorrhizal symbiosis in acid soil condition. Mycorrhizal colonization of *D. glomerata* strongly affected by soil pH, but the *A. odoratum* was not yet, mycorrhizal colonization of *D. glomerata* significantly decreased by declined the soil pH, result to decrease the nutrient uptake, growth, acidity tolerance and competition ability of *D. glomerata*, conducted its declined. On the contrary, mycorrhizal colonization of *A. odoratum* not decreased by declined the soil pH, the mycorrhizal symbiosis led to a positive effect on nutrient uptake, growth, acid tolerance and competition ability of *A. odoratum*. Consider to *A. odoratum* invade and extend in severity acidic artificial pasture origin of this specificity on arbuscular mycorrhizal symbiosis in acidic soil other than *D. glomerata*. Our results have a important meaning and value reference on understand to mechanism of invade and extend in severity acidic artificial pasture by *A. odoratum* and declined of *D. glomerata* in same condition.

References

- Wu En, G.R. Xin, K. Sugawara, et al. (2006). Effect of acidic soil on arbuscular mycorrhizal symbiosis of *Anthoxanthum odoratum* L. *Acta agrestia sinica* 14 (2): 152-155.
- Fitter, A.H. and Nichols, R. (1988). The use of benomyl to control infection by vesicular-arbuscular mycorrhizal fungi. *New phytol.* 110: 201-206.
- Giovannetti M. and Mosse B. (1980). An evaluation of techniques for measuring VAM infection in roots. *New phytol.* 84: 489-500.