



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Growth Characteristic, Productivity and Nutritive Value of Forages by Reseeding Methods in Forest Fire Burnt Pasture

J. K. Lee

Rural Development Administration, South Korea

S. H. Yoon

Rural Development Administration, South Korea

Y. C. Lim

Rural Development Administration, South Korea

H. C. Ji

Rural Development Administration, South Korea

B. S. Jeon

Rural Development Administration, South Korea

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/6-1/30>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

J. K. Lee, S. H. Yoon, Y. C. Lim, H. C. Ji, B. S. Jeon, K. Y. Kim, K. J. Choi, H. S. Park, and S. Seo

Growth characteristic , productivity and nutritive value of forages by reseeding methods in forest fire burnt pasture

J . K . Lee , S . H . Yoon , Y . C . Lim , H . C . Ji , B . S . Jeon , K . Y . Kim , K . J . Choi , H . S . Park and S . Seo
Grassland & Forages Division , National Institute of Animal Science , RDA , Cheonan , Chungnam 330-801 , Korea ,
E-mail : leejkhs@hanmail.net , National Institute of Subtropical Agriculture , RDA , Jeju 690-150 , Korea

Key words : Forest fire , pasture , reseeding , dry matter yield , nutritive value

Introduction Periodic burning is a management tool used to rejuvenate old stands of grasses . Burning maintains the vegetation in a more youthful and productive state for grazing . How a plant responds to fire depends on the height above ground of its growing points , a function of plant maturity and plant-growth characteristics . Limited information is known about the use of fire on cool-season grasses . However , these may include many of the same objectives associated with burning , such as woody plant control or thatch removal . Also , grazing distribution and uniformity may be improved by burning cool-season bunch grasses , which accumulates dead stems that block access to new growth . From April 4-6 , 2005 , there was a huge fire in Yangyang-gun , Kangwon-do , Korea , which burned over 250 ha of forests and 246 buildings including 160 houses . Therefore , effects of reseeding rate and timing on growth characteristic , productivity and nutritive value of forages in the forest fire burnt pasture were investigated in this study .

Materials and methods This study was conducted from April 2005 to December 2006 in a private farm in Kangwon-do , Korea . Six treatments were considered namely : existing vegetation , damaged vegetation (Forest fire burnt pasture in the region) without any modification and damaged vegetation with modifications such as 30% and 50% reseeding 10 days and 20 days after the forest fire . The existing vegetation was seeded with 15 kg of orchardgrass , 10 kg of tall fescue , 3 kg of Kentucky blue grass and 2 kg of white clover . During the establishment period , fertilizer was applied at a rate of 80 kg N , 200 kg P and 70 kg K/ha and 210 kg N , 150 kg P and 180 kg K/ha for management distributed equally across the experimental area in the spring and time after every cutting . The forages were harvested by cuttings 3 times in 2005 and 4 times in 2006 . Estimates of yield were determined by harvesting forage in a 1 m² area in each plot . Fresh forages were oven dried for 72h at 75°C , weighed and converted to DM yield . Forage nutritive value was evaluated in terms of ADF and NDF (Goering & Van Soest , 1970) , and *in vitro* dry matter digestibility (IVDMD ; Moore , 1970) .

Results The DM yield for existing vegetation was the highest in the 1st year(2005) of the study(Table 1) . In the 2nd year study , DM yield obtained from damaged vegetation with no treatment was the highest . The highest yield was not significantly different , however , from the DM yield of other treatments in this study .

Table 1 Dry matter (DM) yield of forages in 2005 and 2006 .

Treatments	DM yield (kg/ha)		
	2005	2006	Mean
Existing vegetation	8 ,587	7 ,803	8 ,195
Forest fire burnt pasture	6 ,296	9 ,482	7 ,889
30% Reseeding , 10days after burning	7 ,045	7 ,772	7 ,408
50% Reseeding , 10days after burning	7 ,910	8 ,288	8 ,099
30% Reseeding , 20days after burning	7 ,332	8 ,049	7 ,690
50% Reseeding , 20days after burning	7 ,100	8 ,879	7 ,989
LSD (0 .05)	NS	NS	NS

Conclusions Although there were no significant yield differences obtained in this study , the mean DM yield for existing vegetation was the highest . A benefit of reseeding burnt pastures was not found . We concluded that forest fire resulted in reduction of the pasture yield for a short period but it did not have any significant effect in the long term .

References

Goering , H . L . , Van Soest , P . J . , (1970) . Forage fiber analysis . *Aggr . Handbook* . No . 379 . USDA .
Moore , J . E . , (1970) . *In vitro* dry matter yield or organic matter digestion . *Nutri . Res . Techn .* 1 :5001-5005 .