



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Aspects of Range Condition Recovery in the Southern Jebel al Akhdar, Northeastern Libya

Adel Mahmoud

Omaral Mukhtar University, Libya

Ashraf Gadallah

Omaral Mukhtar University, Libya

Sanose Mohemmed

Omaral Mukhtar University, Libya

Musbah Mohamed

Garyounis University, Libya

Adel Abdel-Ghani

Omaral Mukhtar University, Libya

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/6-1/34>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

Adel Mahmoud, Ashraf Gadallah, Sanose Mohemmed, Musbah Mohamed, Adel Abdel-Ghani, Ramadan Alhendawi, and Peter J. Russell

Aspects of range condition recovery in the southern Jebel al Akhdar , northeastern Libya

A del Mahmoud¹ , Ashraf Gadallah¹ , Sanose Mohemmed¹ , Musbah Mohamed² , A del Abdel-Ghani¹ , Ramadan Alhendawi¹ and Peter J Russell³

¹ Omar al Mukhtar University , Al Baida , Libya ; ² Garyounis University , Benghazi , Libya ; ³ Curtin University , Locked Bag 22 , Kalgoorlie , Western Australia and corresponding author ; E-mail : p . russell@cmae . curtin . edu . au

Key words : monitoring , enclosure , grazing , soil-condition , vegetation-condition , foliage biomass

Introduction The extensive grazing of sheep , goats and camel occurs in the southern Jebel al Akhdar rangelands of northeast Libya . Climate is Mediterranean , characterised by winter precipitation between October and April . Annual average rainfall declines rapidly from 250 mm in the northern uplands to less than 30 mm in the south . Vegetation is zoned dwarf shrub steppe . Perennial shrubs include *Artemisia herba-alba* , *Haloxylon scoparium* , *Anabasis articulata* , *Suaeda pruinosa* , *S . vera* and *Salsola tetrandra* (SWECO 1986) . Inappropriate and unsustainable historical and contemporary land use practices have resulted in widespread severely degraded soil and vegetation (SWECO 1986 ; Anon . 2004 ; Russell 2006) . The study objective is to quantify range condition improvement in areas protected from grazing .

Methods and materials This integrated study utilised plant inventory surveys and a suite of permanent monitoring sites located within and outside fenced enclosures at three localities to investigate differences in perennial plant species richness , size and density , foliage biomass , energy and nutrient yield , landscape function , soil surface condition and soil micro-nutrients . Plant inventorying was undertaken during the 2006 spring flowering season and the other data were collected in early summer with a second biomass sampling in early autumn 2007 . Monitoring site layout and data collection protocol is based on the Western Australian Rangeland Monitoring System (Watson , Novelly & Thomas 2007) .

Results and discussion Statistically significant differences between enclosures and open areas were found for all vegetation and soil attributes except soil micro-nutrients and some physical and chemical properties . Levels of erosion are less and species richness and plant biomass are markedly higher within enclosures . For example , within the most productive enclosure (Maduar Zetun , natural regrowth protected for six years) , foliage biomass was approximately 222 kg (DM) per hectare and available gross energy was approximately 2 ,182 MJ per hectare , compared to values of 122 and 1 ,198 respectively , in the adjacent open area . A total of 33 perennial species were recorded in the enclosure compared to 22 in the grazed area . Mean soil surface condition (index value 103) , driven mainly by improved water infiltration and soil stability , had improved 27% during the six years of protection compared to the adjacent area subject to intensive , continuous grazing . This study highlights the interaction between vegetation and soil in land degradation . Protection from high grazing pressure allowed vegetation cover , richness and standing biomass to increase with concomitant reduction in soil erosion . Improved soil surface condition probably facilitated plant recruitment .

Conclusions This quantitative study supports visual observations that improvement has occurred in the protected areas but not in unprotected grazed areas . The study also supports forage biomass energy yield improvement predictions made by SWECO (1986) as part of their range management recommendations to protect areas from grazing . It is estimated that at least 10 years of protection in the higher rainfall areas , and longer in the low rainfall areas , is required for range condition to improve sufficiently before being opened to controlled grazing .

References

- Anon . 2004 , *A Reconnaissance Evaluation Study of the Status of Natural Vegetation Cover of Al Jabal Al Akhdar* , Omar al Mukhtar University , Al Baida , Libya .
- Russell , P . 2006 , Range science and desertification in Jebel al Akhdar , *World Environment Day Seminar Series , Faculty of Natural Resources and Environmental Sciences , Omar Mukhtar University , Al Baida , Libya* , p . 3 .
- SWECO 1986 , *Land Survey , Mapping and Pasture Survey for 550 ,000 Hectares of South Jebel el Akhdar Area* , for Socialist People's Libyan Arab Jamahiriya Secretariat for Agricultural Reclamation and Land Development , Contract No 15/90/81 .
- Watson , I . W . , Novelly , P . E . & Thomas , P . W . E . 2007 , Monitoring changes in pastoral rangelands—the Western Australian Rangeland Monitoring System (WARMS) , *The Rangeland Journal* , vol . 29 , no . 2 , pp . 191—205 .