



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Seedling Recruitment in Steppe Communities

Na Zhao

*China Agricultural University, China*

Kun Wang

*China Agricultural University, China*

Zengwen Wang

*Chinese Academy of Sciences, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/6-1/46>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Seedling recruitment in steppe communities

Zhao Na<sup>1</sup>, Wang kun<sup>1</sup>, Wang Zeng wen<sup>2</sup>

<sup>1</sup> China Agricultural University 2 Institute of Botany, Chinese Academy of Sciences, Beijing, 100093 China. E-mail: zhaa88na@163.com

**Key words :** seedling ; density ; seasonal dynamics ; vegetative recruitment ; steppe communities

**Introduction** Seedlings are central to population maintenance, community structure and succession. The conditions of seedling growth can play an important role in local-scale species richness and productivity (Zeiter et al., 2006). Relatively few studies have addressed seedling-recruitment in semiarid grassland ecosystems of Inner Mongolia. The dynamics of seedling recruitment after different types of disturbance is unclear for many species in this region. The aim of the present study was to evaluate the seasonal dynamics of seedlings in typical steppe communities.

**Materials and methods** Three perennial graminoid species from the steppe community of Inner Mongolia were selected for population and seedling studies. The experimental site was located in an area of flat terrain in the southern part of Duolun County, Inner Mongolia. Treatment included mowing and grazing. This experiment was started in Mid-May, and extended to early October. To facilitate seedling monitoring, eight 25cm 25cm quadrates were sampled in each of 5 replicated subplots.

**Result and discussions** Seedling densities of all species showed an increasing trend during the growth period. Seasonal seedling dynamics of *S. krylovii* was similar to that of the Gramineae, which account for majority of the seedlings.




Figure 3 Seasonal dynamics of seedling density.

In a comparison of all three species, the pattern of relative seedling density was  $S. krylovii > C. squarrosa > A. cristatum$  ( $P < 0.05$ ) under control conditions, but under both grazing and mowing treatments the rank order was  $C. squarrosa > S. krylovii > A. cristatum$  ( $P < 0.05$ ). Mowing disturbance results in a reduction of reproductive parts, especially for late-flowering plants. This land use practice, therefore, can induce seed limitations for a species normally producing a sufficient amount of seeds (Overbeck et al., 2003). The species in this experiment, however, all had later reproductive phenology. The tall-growing species- *S. krylovii* lost a larger proportion of biomass than the smaller plants when mowed.

**Conclusions** From the standpoint of seedling density, *S. krylovii* accounted for the majority of all seedlings present in the study plots. It's trend of seedling recruitment, however, was similar to that of the other species. In this typical steppe community, seedlings, especially of the dominant species, played a vital role in population recruitment.

### References

- Overbeck G, Kiehl K, Abs C (2003). Seedling recruitment of *Succisa inflata* in fen meadows: Importance of seed and microsite availability. *Applied Vegetation Science*, 6, 97-104.
- Zeiter M, Stampfli A, Newbery DM (2006). Recruitment limitation constrains local species richness and productivity in dry grassland. *Ecology*, 87, 942-951.