



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

The Effects of Stocking Rate and Corn Gluten Supplemental Feed on the Performance of Young Beef Cows

A. D. Nkengafac
University of Buea, Cameroon

P. Tinta
University of Buea, Cameroon

G. Ayuk
University of Buea, Cameroon

M. Mansue
University of Buea, Cameroon

F. Fuh
University of Buea, Cameroon

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/7-1/36>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

The effects of stocking rate and corn gluten supplemental feed on the performance of young beef cows

A .D .Nkengafac , P .Tinta , G Ayuk , M Mansue and F .Fuh .

Department of Animal Health , Faculty of Science , University of Buea , P .O .Box 63 Buea , Cameroon .E-mail : nkengafac@yahoo .com

Key words : Beef cow , stockpiled forage , corn gluten feed

Aim A winter grazing experiment was conducted to evaluate the effects of stocking rate and corn gluten feed supplementation on forage mass and composition and the BW and BCS of bred 2-yr-old cows grazing stockpiled forage during winter .

Methods Two 12 2-ha blocks containing Fawn , endophyte-free , tall fescue and red clover were each divided into 4 pastures of 2 .53 or 3 .54 ha .Hay was harvested from the pastures in June and August of 2005 and 2006 , and N was applied at 50 .5 kg/ha at the initiation of stockpiling in August .On October 22 , 2005 , and October 20 , 2006 , twenty-four 30-mo-old Angus-Simmental and Angus cows were allotted by BW and BCS to strip-graze for 147 d at 0 .84 or 1 .19 cow/ha .Eight similar cows were allotted to 2 dry lots and fed tall fescue-red clover hay *ad libitum* .Corn gluten feed was fed to cows in 2 pastures to maintain a mean BCS of 5 (9-point scale) at each stocking rate and in the dry lots (high supplementation level) or when weather prevented grazing (low supplementation level) in the remaining 2 pastures at each stocking rate .

Results Mean concentrations of CP in yr 1 and 2 and IVDMD in yr 2 were greater ($P < 0 .10$) in hay than stockpiled forage over the winter .At the end of grazing , cows fed hay in dry lots had greater ($P < 0 .05$) BCS in yr 1 and greater ($P < 0 .10$) BW in yr 2 than grazing cows .Grazing cows in the high supplementation treatment had greater ($P < 0 .10$) BW than cows grazing at the low supplementation level in yr 1 .Cows in the dry lots were fed 2 ,565 and 2 ,158 kg of hay DM/cow .Amounts of corn gluten feed supplemented to cows in yr 1 and 2 were 46 and 60 kg/ cow and did not differ ($P = 0 .33$, yr 1 ; $P = 0 .50$, yr 2) between cows fed hay or grazing stockpiled forage in either year .

Conclusion Estimated production costs were greater for cows in the dry lots because of hay feeding .