



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII  
International Rangeland Congress

---

## Carbon Balance in Stand of an Annual Herb at an Elevated CO<sub>2</sub> Concentration

Almaz Borjigidai  
*Central University for Nationalities, China*

Kouki Hikosaka  
*Tohoku University, Japan*

Tadaki Hirose  
*Tohoku University, Japan*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/8-1/4>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Carbon balance in stand of an annual herb at an elevated CO<sub>2</sub> concentration

Almaz Borjigidail<sup>1,2\*</sup>, Kouki Hikosaka<sup>2</sup> and Tadaki Hirose<sup>2,3</sup>

<sup>1</sup> Central University for Nationalities, China.


<sup>2</sup> Tohoku University, Japan.

<sup>3</sup> Tokyo University of Agriculture, Japan. E-mail: [almaz\\_b@hotmail.com](mailto:almaz_b@hotmail.com)


**Key words :** carbon balance, stand, photosynthesis, respiration, elevated CO<sub>2</sub> concentration, *Chenopodium album*

**Introduction** Global atmospheric CO<sub>2</sub> concentration is predicted to double by the end of the present century (IPCC, 2001). The influence of increased CO<sub>2</sub> on plant and ecosystem functions has been a major concern in ecology and global change sciences (Long et al., 2004). In most studies, elevated CO<sub>2</sub> enhanced canopy photosynthesis. However, the magnitude of the enhancement varies among studies (Nowak et al., 2004). More recently, Sakai et al. (2006) showed that enhancement of canopy carbon gain by elevated CO<sub>2</sub> is sensitive to the growth stage and leaf nitrogen concentration in the rice stand. However, since most of these studies determined photosynthesis in the stand without destructive harvests, they were not able to fully analyze mechanisms involved in variation in stand carbon balance. Canopy photosynthesis is a function of canopy structure (leaf area index, K, leaf nitrogen distribution etc.), leaf physiology (photosynthesis, respiration etc.) and environmental factors (irradiance, N availability, temperature etc.). In the present study, we studied the effect of elevated CO<sub>2</sub> on carbon balance for stands of an annual *Chenopodium album*.

**Materials and methods** Stands of *C. album* were established in open top chambers at ambient and elevated CO<sub>2</sub> concentrations (370 and 700 μmol mol<sup>-1</sup>) in the experimental garden of Tohoku University, Sendai, Japan (38°15'N, 140°52'E). Plant dry mass growth, photosynthesis and respiration were determined through the growing season. CO<sub>2</sub> exchange of the stand was estimated with a canopy photosynthesis model (Hirose, 2005). Rates of light-saturated photosynthesis (Pmax) and dark respiration (Rm) of leaves as related with nitrogen content per unit leaf area (Narea) and time-dependent reduction in specific respiration rates (SRR) of stems and roots were incorporated into the model.



**Figure 1** Comparison of stand carbon balance (canopy photosynthesis minus stems and roots respiration) with mean daily plant growth rates (GR) in *C. album* stands. Carbon balance is a mean of calculated values for a period between harvests (29 and 41, 42 and 62, 63 and 81 DAT). Open circles, ambient CO<sub>2</sub> (370 μmol mol<sup>-1</sup>); closed circles, elevated CO<sub>2</sub> (700 μmol mol<sup>-1</sup>). Solid line represents regression line:  $y = 0.034x$ ,  $r = 0.99$ . Dotted line indicates equivalence of 1 mol CO<sub>2</sub> with 30 g dry mass.



**Figure 2** Enhancement of canopy photosynthesis by elevated CO<sub>2</sub> as a function of DAT (Days after transplanting). Each data point expresses a mean of canopy photosynthesis. Results of sensitivity analyses (Photosynthesis-effect, triangles, and Structure effect, squares) are also shown.

**Results and conclusions** Daily canopy carbon balance, calculated as an integration of leaf photosynthesis minus stem and root respiration, showed a fairly good agreement with dry mass growth determined by harvests (Figure 1). The enhancement of canopy photosynthesis with elevated CO<sub>2</sub> was 80% at an early stage and decreased to 55% at flowering. Sensitivity analyses suggested that an alteration in leaf photosynthetic traits enhanced canopy photosynthesis by 40-60% throughout the experiment period, whereas altered canopy structure (leaf area index, leaf nitrogen distribution and light gradient in the canopy) contributed to the enhancement at the early stage only (Figure 2). In early stages when plant size was small, nutrient supply might not be a limiting factor of plant growth, leading to an enhancement of LAI at elevated CO<sub>2</sub>, while in later stages of plant growth, nutrient supply became more limiting and the structural difference might diminish between the two growth CO<sub>2</sub> concentrations. Thus various factors interact to determine

the stand carbon balance that is influenced by elevated CO<sub>2</sub>.

### References

- Hirose T (2005) Development of the Monsi-Saeki theory on canopy structure and function. *Annals of Botany* 95 : 483-494 .  
 Long SP, Ainsworth EA, Rogers A, Ort DR (2004) Rising atmospheric carbon dioxide: Plants FACE the Future. *Annual Review of Plant Biology* 55 : 591-628 .  
 Nowak R, Ellsworth D, Smith S (2004) Functional responses of plants to elevated atmospheric CO<sub>2</sub>-do photosynthetic and productivity data from FACE experiments support early predictions? *New Phytologist* 162 : 253-280 .  
 Sakai H, Hasegawa T, Kobayashi K (2006) Enhancement of rice canopy carbon gain by elevated CO<sub>2</sub> is sensitive to growth stage and leaf nitrogen concentration. *New Phytologist* 170 : 321-332 .