



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Climate Change and Human Activity Impacts on the Net Primary Production of Alpine Grassland in Northern Tibet, China

Qingzhu Gao

*Chinese Academy of Agricultural Sciences, China*

Yue Li

*Chinese Academy of Agricultural Sciences, China*

Yunfan Wan

*Chinese Academy of Agricultural Sciences, China*

Wangzha Jiangcun

*Nagqu Bureau of Agriculture and Husbandry Management Department, China*

Baoshan Wang

*Nagqu Bureau of Agriculture and Husbandry Management Department, China*

*See next page for additional authors*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/8-1/15>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

**Presenter Information**

Qingzhu Gao, Yue Li, Yunfan Wan, Wangzha Jiangcun, Baoshan Wang, and Wenfu Li

## Climate change and human activity impacts on the net primary production of alpine grassland in northern Tibet , China

Gao Qingzhu<sup>1</sup> , Li Yue<sup>1</sup> , Wan Yunfan<sup>1</sup> , Jiangcun Wangzha<sup>2</sup> , Wang Baoshan<sup>2</sup> , Li Wenfu<sup>2</sup>


<sup>1</sup> Institute of Environment and Sustainable Development in Agriculture , CAAS , Beijing 100081 , E-mail :gaoqzh@ami.ac.cn , <sup>2</sup> Nagqu Bureau of Agriculture and Husbandry Management Department , Tibet Autonomous Region , Nagqu , 852100

**Key words :** NPP , climate change , human activity , trends analysis , impact assessment , alpine grassland , northern Tibet

**Introduction** Northern Tibet region is the headstreams of major rivers , including Yangtze River , Nujiang River , and Lancangjiang River in China (Gao et al , 2005) . Based on the remote sensing data from 1981 to 2004 as well as other related data , spatial trend of alpine grassland net primary production (NPP) and its resonances to climate change and human activity in Northern Tibet were analyzed with the help of geographical information system .

**Material and methods** Alpine grassland NPP were calculated by the CASA (Carnegie-Ames-Stanford Approach) model and validated by comparing with observational data . Pearson coefficient was used to predict the long term variations of alpine grassland NPP from 1981 to 2004 . A correlative analysis was made to calculate spatial correlation coefficients between alpine grassland NPP and climate change in Northern Tibet . The impacts of human activity intensity was analyzed on alpine grassland NPP based on GIS techniques .

**Results** The change in alpine grassland in most areas of Northern Tibet was not obvious across roughly 89% of total grassland area ; while the area with marked change only accounts for 11.4% , with roughly 11.3% showing decrease and < 0.1% increase (Figure 1) . In recent years , the precipitation variation in Northern Tibet resulted in an increase of grassland NPP , though solar radiation resulted in decreased grassland NPP . During the period of 1981-2004 , climate factors affected the grassland NPP in Northern Tibet in the following orders : total solar radiation > precipitation > temperature (Figure 1) . The negative effects of local residential areas on the rate of grassland NPP change are smaller than that of roads . In general the intensity of human activity in the region near to road and the residential area are strong and the influence on grassland NPP change tendency are bigger .



(a) Trends of NPP ; (b) NPP and precipitation ; (c) NPP and temperature ; (d) NPP and solar radiation  
ESD is extremely significant decrease ; SD is significant decrease ; ISD is insignificant decrease ; ISI is insignificant increase ; SI is significant increase ; ESI is extremely significant increase ; ESNC is extremely significant negative correlation ; SNC is significant negative correlation ; NCNS is negative correlation but none significant ; PCNS is positive correlation but none significant ; SPC is significant positive correlation ; ESI is extremely significant positive correlation

**Figure 1** Trends of grassland NPP and spatial correlation between alpine grassland NPP and annual precipitation , annual mean temperature and annual solar radiation in Northern Tibet .

**Conclusions** Most areas in Northern Tibet did not show a significant annual NPP change . The negative effects of local residential areas on the rate of grassland NPP change are smaller than that of roads . During the period of 1981-2004 , climate factors affected the grassland NPP in the following orders : total solar radiation > precipitation > temperature . Generally , the impact of regional climate change on grassland NPP was more negative than positive .

### References

- Gao Qingzhu , Li Yue , Lin Erda , 2005 . Temporal and spatial distribution of grassland degradation in Northern Tibet , *Acta Geographica Sinica* , 60(6) :965-973 .  
Nemani R.R , Keeling C.D , Hashimoto H . 2003 . Climate-driven increases in global terrestrial net primary production from 1982 to 1999 . *Science* , 300 :1560-1563 .