



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Study on Methane Diurnal Emission nearby Cuona Lake in Naqu of Tibet with Different Humidity

Zhongxin Zhang

*Capital Normal University, China*

Yunfan Wan

*Chinese Academy of Agricultural Sciences, China*

Yu'e Li

*Capital Normal University, China*

Qingzhu Gao

*Chinese Academy of Agricultural Sciences, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/8-1/46>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Study on methane diurnal emission nearby Cuona Lake in Naqu of Tibet with different humidity

Zhang Zhong-xin<sup>1,2</sup>, Wan Yun-fan<sup>2</sup>, Li Yu-e<sup>1,2</sup>, Gao Qing-zhu<sup>2</sup>

<sup>1</sup>College of Resource Environment and Tourism; Capital Normal University; Beijing 100037; China, E-mail: zhangzhx@ami.ac.cn, <sup>2</sup>Institute of Environment and Sustainable Development in Agriculture; Chinese Academy of Agricultural Sciences; Beijing 100081

**Key words** methane diurnal emission, Cuona lake, wetland, humidity, Naqu, Tibet

**Introduction** It shows that the 70~80 percentage of atmospheric CH<sub>4</sub> is from biogenic sources. And it is believed that paddy fields and wetlands are the dominant sources of atmospheric CH<sub>4</sub> (Hu, 2005). We made a study on methane emission from the wetland nearby Cuona Lake in Naqu of Tibet with different humidity by field investigation, which can contribute to evaluating methane emission from wetlands in Tibet.

**Materials and methods** The field experiment site was located in nearby Cuona Lake in Naqu of Tibet. On August 2006, 3 treatments of CH<sub>4</sub> flux were imposed with 3 replicates according to the distance from the lake. Treatments were: (1) Lake level treatment (LT); (2) Wet treatment (WT), 100 meter far from the lake; (3) Dry treatment (DT), 500 meter far from the lake. LH, WH and DH were the humidity of 3 treatments, respectively. Using the enclosed chamber technique, CH<sub>4</sub> fluxes were measured. The chamber was a cylinder of 40 centimeter high and its base diameter was 30 centimeters. When measured, the base was inserted 5-centimeter underground, and sealed by distilled water. Samples collected by syringe were transferred into plastic gas bags for measurement by HP6890GC.

**Results** By the curve, it was proved that LH, WH and DH gradually decreased with the distance from lake. But it wasn't significant for all of treatments diurnal humidity variation. CH<sub>4</sub> flux of all treatments with time was as follows: WT > DT > LT. The trend of CH<sub>4</sub> diurnal flux of all treatments was accord with each other. A peak appeared in 3 curves respectively. WT peak and DT peak appeared at 15 pm, and LT peak was at 17 pm.




Figure 1 Diurnal CH<sub>4</sub> flux and humidity dynamic nearby Cuona lake in Naqu of Tibet.

**Conclusions** CH<sub>4</sub> emission flux of all treatments was positive, which shown that this wetland was the source of atmospheric CH<sub>4</sub>. The CH<sub>4</sub> flux of LT was smaller, which may result in the inhabitation of CH<sub>4</sub> emission due to excessive humidity of the lake-level. Because of the far distance between DT and lake and smaller humidity, it didn't result in much higher CH<sub>4</sub> flux. With the moderate distance between WT and lake and with appropriate humidity, it could accelerate CH<sub>4</sub> emission.

### Reference

Hu Qiwu, Wu Qin, Li Dong, Cao Guangmin, 2005. Comparative study on methane emissions from alpine grasslands with different soil water content. *Chinese Journal of Ecology* 24 (2), 118-122.