



## Rotational and Continuous Grazing of Sheep in the Inner Mongolian Steppe of China

C. J. Wang

*Christian-Albrechts-University Kiel, Germany*

B. M. Tas

*Christian-Albrechts-University Kiel, Germany*

T. Glindemann

*Christian-Albrechts-University Kiel, Germany*

K. Mueller

*Christian-Albrechts-University Kiel, Germany*

A. Schiborra

*Christian-Albrechts-University Kiel, Germany*

*See next page for additional authors*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/9-3/19>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

### **Presenter Information**

C. J. Wang, B. M. Tas, T. Glindemann, K. Mueller, A. Schiborra, P. Schoenbach, M. Gierus, Friedhelm Taube, and A. Susenbeth

## Rotational and continuous grazing of sheep in the Inner Mongolian steppe of China

C. J. Wang, B. M. Tas, T. Glindemann, K. Mueller, A. Schiborra, P. Schoenbach, M. Gierus, F. Taube and A. Susenbeth  
 Institute of Animal Nutrition and Physiology, Christian-Albrechts-University Kiel, Hermann-Rodewald-Str. 9, 24118 Kiel, Germany, E-mail address: susenbeth@aninut.uni-kiel.de

**Key words** : rotational and continuous grazing, feed intake, digestibility, live weight gain

**Introduction** The Inner Mongolia steppe of China is traditionally used for grazing. In the last three decades overgrazing by sheep led to a sparse cover of vegetation in winter that enables soil erosion and hence degradation of the grassland, which may be influenced by grazing system. The objective of this study—which is part of the Sino-German DFG research cooperation MAGIM—was to investigate the effects of rotational and continuous grazing on chemical composition of herbage, herbage mass (HM), feed intake and digestibility and live weight gain (LWG) of sheep grazing at a moderate stocking rate of 4.5 sheep/ha on the Inner Mongolian steppe of China.

**Materials and methods** In the growing season from mid June until mid September in 2005 and 2006 a grazing experiment was conducted comparing a continuous and rotational grazing system with one replication. In the continuous system the sheep remained on one 2 ha plot throughout the season, whereas in the rotational system the plot was subdivided into four paddocks of 0.5 ha each and sheep were moved to the next paddock every 10 days. The stocking rate was 4.5 yearling sheep per ha. The chemical composition of herbage was determined with NIRS and HM was measured using enclosure cages. The intake of organic matter (OMI) was calculated by daily output of fecal OM, estimated by the external marker titanium dioxide, and by OM digestibility (dOM) of herbage ingested by sheep. The dOM was estimated by the fecal crude protein (CP) method (Wang 2007). Intake of digestible OM (dOMI) was calculated by OMI multiplying with dOM, and intake of metabolisable energy (MEI, MJ/d) by multiplying dOMI (kg/d) with 16 (MJ/kg), as described by McDonald et al. (1988).

**Results** The concentration of CP was lower and NDF higher with rotational than with continuous grazing. The OMI, dOM and MEI with rotational grazing were lower with 1.09 kg/d, 54.9% and 9.6 MJ/d, respectively, than with continuous grazing with 1.31 kg/d, 57.4% and 12.0 MJ/d, respectively. However, the grazing systems did not affect HM and LWG of sheep. The annual precipitation was low with 166 mm in 2005 and 304 mm in 2006 compared with the average value of the last 30 years of 350 mm, which influenced the measured variables in different years.

**Table 1** The measured variables in rotational and continuous grazing system in 2005 and 2006.

Item	Rotational grazing		Continuous grazing		SEM	P value	
	2005	2006	2005	2006		GS	Year
OM %	94.3	93.1	94.9	92.7	0.12	0.78	<0.001
CP %	8.1	11.9	8.8	13.0	0.21	0.001	<0.001
NDF %	73.3	69.1	72.0	67.8	0.24	0.016	<0.001
HM, kg/ha	592	490	546	314	35.9	0.14	0.03
OMI, kg/d/sheep	1.03	1.14	1.21	1.27	0.02	<0.001	<0.001
dOM %	53.3	56.5	56.3	58.4	0.14	<0.001	<0.001
MEI, MJ/d/sheep	8.8	10.3	10.9	11.3	0.17	<0.001	<0.001
LWG, g/day	68	84	62	81	2.53	0.41	0.01

**Conclusions** The results based on these two years showed that the grazing system affected quality of feed, digestibility and intake in sheep grazing on the Inner Mongolian steppe, but the LWG was not affected. Further studies are required in years with mean precipitations before final evaluation of the grazing systems can be made.

### References

- Wang CJ (2007) Dissertation, University of Kiel.  
 McDonald P, Edwards RA, Greenhalgh JFD (1988) *Animal Nutrition*. Copublished in United State with John Wiley and Sons, Inc., New York. pp. 245-259.