



Characterization of Soil Properties under Legume and Non-Legume Tree Canopies Occurring in Signal Grass (*Brachiaria decumbens* Stapf.) Pastures

José C. B. Dubeux Jr.

Universidade Federal Rural de Pernambuco, Brazil

M. A. Lira

Instituto Agronômico de Pernambuco, Brazil

M. V. F. Santos

Universidade Federal Rural de Pernambuco, Brazil

D. M. Barros

Universidade Federal Rural de Pernambuco, Brazil

M. A. Silva

Universidade Federal Rural de Pernambuco, Brazil

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/10-1/14>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

Presenter Information

José C. B. Dubeux Jr., M. A. Lira, M. V. F. Santos, D. M. Barros, M. A. Silva, and Alexandre C. L. de Mello

Characterization of soil properties under legume and non-legume tree canopies occurring in signal grass (*Brachiaria decumbens* Stapf.) pastures

J.C.B. Dubeux, Jr.¹; M.A. Lira²; M.V.F. Santos¹; D.M. Barros¹; M.A. Silva¹; A.C.L. Mello¹

¹UFRPE-Departamento de Zootecnia, R. Dom Manoel de Medeiros, S/N, Dois Irmaos, Recife, PE, Brazil, 52171-900; e-mail: dubeux@dz.ufrpe.br; ²IPA-Av. General San Martin, Bongi, Recife, PE, Brazil.

Key words: agroforestry, nutrient cycling, shade, soil fertility, soil organic matter

Introduction Trees improve the soil by numerous processes, mainly when used in agroforestry systems, where different species are cultivated in the same area. Legume trees influence nutrient availability within the root zone of the intercropped species by different processes such as: i) N biologic fixation; ii) nutrient recovery from deeper soil layers; iii) reduction in soil erosion and runoff; and iv) increasing nutrient bioavailability by increasing soil organic matter (SOM) mineralization. The objective of this research was to evaluate the total soil organic matter and soil Mehlich-I P at different points underneath the tree crown, according to a shade gradient of legume and non-legume trees occurring on signalgrass (*Brachiaria decumbens* Stapf.) pastures.

Material and methods The experiment was carried out in the coastal region of Pernambuco State, in Itambe Agricultural Research Station from IPA. Geographical coordinates are 7°25'0"S lat. and 35°6'0"W longitude with 190 m above sea level. Average annual rainfall in this area is 1,300 mm and the soil is classified as an oxisol. Soil samples were taken according to the crown projection of two legume trees (*Mimosa caesalpiniiifolia* Benth. and *Machaerium aculeatum* raddi.) and two non-legume trees (*Anacardium occidentale* L. and *Artocarpus integrifolia* L.). Samples were taken at 10%, 50%, 100%, and 150% of shade projection at the 0-10 cm soil layer. Soil analyses of these samples were accomplished at UFRPE, determining total SOM and soil Mehlich-I P (EMBRAPA, 1999). Data was analyzed as complete randomized design (proc mixed SAS) and non-linear and linear models were used (proc nonlin SAS) to fit the data whenever the shade gradient was significant.

Results Soil fertility was greater under the crown projection as shown by SOM (Table 1) and Mehlich-I P (Table 2) for all evaluated tree species. Total SOM showed a negative exponential pattern ($P < 0.01$), with greater SOM concentration closer to the tree trunk. Legume trees showed less SOM accumulation than non-legume trees. This might be related to tree age and crown projection. Non-legume trees had a wider and denser crown than the legume ones. Shade may affect soil temperature and soil microbe activity, modifying decomposition rates. In addition to that, residue deposition likely differs among evaluated species, affecting SOM accumulation. Available P (Mehlich-I) showed a linear decrease, with sun-exposed areas showing less soil P. Considering that these trees occur in a pasture area, cattle manure deposition may contribute to increase soil P under the crown projection (Dubeux et al., 2007).

Table 1 Total SOM (g/kg) according to a shade gradient under legume and non-legume trees occurring on signalgrass pastures, Itambé-PE, Brazil.

Species	Crown projection (%)			
	10	50	100	150
	— g OM/kg soil —			
<i>A. occidentale</i> L.	67.4	53.6	48.8	46.6
<i>M. aculeatum</i> Raddi.	35.9	34.8	28.8	28.6
<i>A. integrifolia</i> L.	73.3	60.9	48.2	41.4
<i>M. caesalpiniiifolia</i> Benth.	35.6	35.2	38.0	30.0
Shade vs. Sun				$P < 0.0002$
Legume tree vs. Non-legume tree				$P < 0.0001$

Standard error = 4.6 g of O.M./kg of soil

Table 2 Mehlich-I P (mg/dm³) according to a shade gradient under legume and non-legume trees occurring on signalgrass pastures, Itambé-PE, Brazil.

Crown projection (%)	Soil Mehlich-I P (mg/dm ³)
10	23.4
50	15.9
100	17.0
150	9.7
S.E.	6.3
Linear effect	$P < 0.0173$
Shade vs. Sun	$P < 0.0350$

Conclusions Soil fertility was greater under the crown projection than under full-sun exposed areas. Greater residue deposition and cattle manure deposition on shaded areas likely contributed to increasing soil fertility levels under trees. Non-legume trees had denser and wider crowns, which likely contributed to greater SOM accumulation under the crowns compared to legume trees.

References

- EMBRAPA-Serviço Nacional de Levantamento e Conservação do Solo, (Rio de Janeiro, R.J.). 1979. *Manual de Métodos de Análise de Solos*, parte 3, análise e fertilidade. Rio de Janeiro, 1979. 17 p.
- Dubeux, J.C.B., Jr.; Sollenberger, L.E.; Mathews, B.W.; Scholberg, J.M.; Santos, H.Q. 2007. Nutrient cycling in warm-climate grasslands. *Crop Sci.*, 47, p.915-928.