



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Lead Hyperaccumulation and Tolerance in *Eremochloa ophiuroides* and *Paspalum vaginatum*

Yanhua Zhu  
*Shanghai Jiaotong University, China*

Zhaolong Wang  
*Shanghai Jiaotong University, China*

Bingru Huang  
*Shanghai Jiaotong University, China*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/11-1/23>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Lead hyperaccumulation and tolerance in *Eremochloa ophiuroides* and *Paspalum vaginatum*

Yanhua Zhu Zhaolong Wang\* and Bingru Huang

School of Agriculture & Biology, Shanghai JiaoTong University, Shanghai 201101, P.R.China, E-mail: turf@sjtu.edu.cn, \* Corresponding author.

**Key words** : *eremochloa ophiuroides*, *Paspalum vaginatum*, lead phytoremediation, phytostabilization

**Introduction** Soil lead (Pb) contamination is one of the more serious environmental problems, with significant impact on human health. The success of phytoremediation for Pb-contaminated soil is mainly dependent on using efficient plants (Pilon-Smits E., 2005). Turfgrasses can effectively control contamination from soil erosion and reduce the chance for human contact with pollutants (Bacon et al., 2005). In the present study, *Eremochloa ophiuroides* and *Paspalum vaginatum* were selected to: (1) examine whether they can tolerate high doses of Pb in contaminated soil; and (2) investigate the efficiency of Pb extraction and accumulation by the two turfgrass species.

**Materials and methods** Two-year turf sod of *E. ophiuroides* and *P. vaginatum* were planted in plastic pots (18 cm in diameter and 15 cm deep) filled with 2.5 kg of a soil mixture contaminated with (5000 mg kg<sup>-1</sup>) and a clean soil control with background Pb concentration of 28.02 mg kg<sup>-1</sup>. Two turfgrass species: *E. ophiuroides* and *P. vaginatum*, were planted in each. The pots were arranged in a randomized complete block design with 4 replicates. Turfgrass canopy photosynthetic rate, leaf chlorophyll content, shoot density, turf quality, shoot and root biomass, and plant Pb concentration were measured.

**Results** Both *E. ophiuroides* and *P. vaginatum* showed excellent tolerance of Pb contamination in the experiment. No significant toxicity symptoms (chlorosis, burning of leaf margins, leaf abscission and shoot dieback) were observed in both grasses. Pb treatment did not show any significant difference in leaf chlorophyll content and shoot density in both grasses. Turf visual quality of *P. vaginatum* did not show any significant difference between Pb treatment and the control, but *E. ophiuroides* showed a decrease of turf visual quality (Table 1). *P. vaginatum* and *E. ophiuroides* accumulated over 7 times and 4 times more Pb concentrations in roots than in the soil when grown in the Pb contaminated soil (Table 2). However, the root:shoot ratio was 16.78% in *P. vaginatum* plant, but 58.13% in *E. ophiuroides*.

**Table 1** Effects of Pb on the growth of turfgrass plants.

Turfgrass species	Treatment	Photosynthetic rate (μmol CO <sub>2</sub> m <sup>-2</sup> s <sup>-1</sup> )	Chlorophyll content (mg g <sup>-1</sup> )	Shoot density (tillers/100 cm <sup>-2</sup> )	Turf visual quality
<i>E. ophiuroides</i>	Control	8.59b <sup>†</sup>	248.8a	51.0a	6.0a
	Pb contaminated soil	12.26a	252.5a	60.3a	4.3b
<i>P. Vaginatum</i>	Control	9.57b	319.0a	161a	8.0a
	Pb contaminated soil	11.32a	313.0a	153a	9.0a

<sup>†</sup> Means followed by the different letters (a, b) were significantly different between Pb contaminated soil and the control within the same species on the basis of LSD test ( $P=0.05$ ).

**Table 2** Pb uptake and distribution in turfgrass plants.

Turfgrass species	Treatment	Pb concentration (mg kg <sup>-1</sup> DW)			Root to shoot transfer rate(%)
		Soil	Root	Shoot	
<i>E. ophiuroides</i>	Control	28±3.5	107±22	72±12.5	
	Pbcontaminated soil	5097±754	18139±1053	10545±38	58.13
<i>P. Vaginatum</i>	Control	28±3.5	68±1.3	27±3.0	
	Pbcontaminated soil	4118±99	29306±116	4918±556	16.78

**Conclusion** *E. ophiuroides* and *P. vaginatum* have good potential for phytoextraction of Pb contaminated soils.