University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress /
VIII International Rangeland Congress

Agrobacterium-Mediated Genetic Transformation of *Elymus breviaristatus* with *Pseudomonas Pseudoalcaligenes* Insecticidal Protein Gene

Daxu Li

Sichuan Academy of Grassland Science, China

Shiqie Bai

Sichuan Academy of Grassland Science, China

Gang Liu

Sichuan Academy of Grassland Science, China

Minghong You

Sichuan Academy of Grassland Science, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/12-1/10>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Agrobacterium-mediated genetic transformation of *Elymus breviaristatus* with pseudomonas pseudoalcaligenes insecticidal protein gene

Li Daxu , Bai Shiqie , Liu Gang , You Minghong

Sichuan Grassland Science Academy , Chengdu , 611731 , China . E-mail : ldx136@163 .com

Key words : Agrobacterium-mediated transformation , callus regeneration , mature embryo , ppip , transgenic plant

Introduction *Elymus breviaristatus* is the predominant type of pasture grass in cold season . This forage grass , however , is a favorable food for grasshoppers (*Locusta Orthoptera*) . Grasshopper plague has become a major damage to the maintenance of pasturage . The development of transgenic technology provides an alternative and effective method for reducing the damage of grasshoppers by the products of pesticidal genes (Akutsu *et al* , 2004) .

Materials and methods Embryogenic calli of *Elymus breviaristatus* were induced from mature seeds , and proliferated on callus induction and maintenance medium containing kinetin 0 .05 mg/L and 2 ,4-D 8 .0 mg/L under dim-light condition at 26°C . The embryogenic calli produced were used for transformation mediated by EHA105 which carries the pCambia 1304-ppip . The calli transformed were selected and grown in the presence of 60 mg/L hygromycin (Li *et al* , 2006) .

Results MS medium was more favorable for callus induction and regeneration of *Elymus breviaristatus* than other media when mature embryos of seeds were used as explants (data not shown) . The calli of three-and four-month-old what had higher frequency of hygromycin-resistant callus when co-cultivated at 19-23°C (data not shown) . The transformation frequency of the calli reached 10 .52 % (data not shown) . Calluses induction and regeneration of *Elymus breviaristatus* are shown in Figure 1 . The presence of *ppip* gene in the genomic DNA of regenerated plants was detected by means of Southern hybridization (Figure 2) , and the expression of the trans-genes was verified by reverse transcription-PCR (Figure 3) .

Figure 1 Calluses induction and regeneration of *Elymus breviaristatus* .

Figure 2 DNA gel blots analysis of transgenic *Elymus breviaristatus* .

Figure 3 RT-PCR analysis of transgenic *Elymus breviaristatus* .

Conclusions *Agrobacterium-mediated* transformation in the conditions tested was effective for transferring foreign genes into *Elymus breviaristatus* . These results may set an example for the *Agrobacterium* method that can effectively transfer foreign genes into *Elymus breviaristatus* .

References

- Akutsu M , Ishizaki T , Sato H (2004) Transformation of the monocotyledonous *Alstroemeria* by *Agrobacterium tumefaciens* . *Plant Cell Rep* . 22(8) : 561-8 .
- Li DX , Zhang J , Zhao J , Zhang Y , Li L , Liu SJ , Chen F , Yang ZR (2006) . Transformation of calli of siberian wildrye grass (*Elymus sibiricus* L . cv . chuancao No .2) mediated by *agrobacterium* . *Journal of Plant Physiology and Molecular Biology* . 2006 , 32 (1) : 45-51 .