



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Identification, Evaluation and Molecular Characterization of *Stylosanthes seabrana*—a Potential and Nutritious Range Legume Having Wider Applicability in India

A. Chandra

Indian Grassland and Fodder Research Institute, India

R. K. Bhatt

Indian Grassland and Fodder Research Institute, India

A. B. Majumdar

Indian Grassland and Fodder Research Institute, India

S. Kumar

Indian Grassland and Fodder Research Institute, India

D. Nagaich

Indian Grassland and Fodder Research Institute, India

See next page for additional authors

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/12-1/32>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Presenter Information

A. Chandra, R. K. Bhatt, A. B. Majumdar, S. Kumar, D. Nagaich, and K. Kumar

Identification , evaluation and molecular characterization of *Stylosanthes seabrana* — a potential and nutritious range legume having wider applicability in India

A . Chandra* , R . K . Bhatt , A . B . Majumdar , S . Kumar , D . Nagaich and K . Kumar
Indian Grassland and Fodder Research Institute , Jhansi-284003 , India

* E-mail of corresponding author : amaresh62@yahoo .com

Key words : drought , sequence tagged sites , specific leaf area , *Stylosanthes*

Introduction The *Stylosanthes* (Fabaceae) genus has wide distribution in tropical , subtropical and temperate regions of the Americas , Africa and South East Asia . In India , *Stylosanthes* is extensively utilized in pastoral , agropastoral and silvipastoral systems (Chandra et al . , 2006) . The characterization of new species in genus *Stylosanthes* is always demanding . The newly identified *S . seabrana* is one such species established well in India without specific rhizobium and is superior in nutritional quality namely essential amino acids . Importance of this species has been also emphasized in other countries like China , Australia and Colombia (Phailkaew et al . , 2004) . It is also better for leaf meal production and a suitable replacement of concentrate for livestock . However , limited numbers of lines in this species pose problems in its wider applications . Here , we report the recovery of ten new lines of *S . seabrana* from existing stocks of *S . scabra* utilizing STS markers which were further supported by morphological attributes . These lines were molecularly characterized using RAPD and STS markers to estimate the level of variability and genetic distance from *S . scabra* of which *S . seabrana* is one of the diploid progenitors (Liu and Musial , 1997) . These lines were also assessed for drought tolerance parameters like specific leaf area , osmolality , proline and soluble protein contents as better drought tolerance lines can be used in artificial synthesis of allo-tetraploid *S . scabra* (Tewari and Chandra , 2008) .

Materials and methods In total 48 accessions procured from ILRI , Ethiopia as *S . scabra* genetic stocks and nine known *S . seabrana* lines from Australia were established at Indian Grassland and Fodder Research Institute , Jhansi and were morphologically evaluated . Specific leaf area (SLA) was measured by randomly harvesting six trifoliate leaves . Osmolality was measured using a vapour pressure osmometer . Proline was estimated colorimetrically . Genomic DNA was isolated using buffer S (Liu and Musial , 1997) . RAPD and STS input binary data matrix was developed by entering 1 to presence or 0 to absence of bands . Dice similarity coefficient was processed for SAHN cluster analysis using UPGMA and tree display was followed for generation of dendrogram (NTSYS tree phenogram) .

Results and discussion Of the 48 accessions of *S . scabra* , 10 distinct lines were identified on the basis of early flowering , leaf shape and less hardness of stem in comparison to *S . scabra* . However , in all these ten lines as in *S . scabra* an axis rudiment , a small appendage at the base of the pod or loment was observed . These characteristics clearly divided these lines from the existing stocks of *S . scabra* . When they were assessed with genome specific marker like sequence - tagged-sites (STS) , all displayed STS phenotypes of typical diploid species . With primer pairs SHST3F3/R3 all putative *S . seabrana* yielded single band of ~550 bp and *S . viscosa* of ~870 bp whereas both of these bands were observed in allo-tetraploid *S . scabra* . Since this primer pairs is known to amplify single or no band with diploid and two bands with tetraploid species , the amplification patterns corroborated that all newly identified *S . seabrana* lines were diploid in nature . Flow-cytometric analysis also indicated the proportional sharing of DNA content between *S . scabra* , and *S . viscosa* and recovered *S . seabrana* lines further indicated the identified lines as *S . seabrana* . Thirty-two RAPD and 17 STS primers indicated more than 92 % similarity between any two accessions of newly identified lines . Low levels of variability in osmolality , SLA , proline and protein content were also observed among the identified *S . seabrana* lines . Results indicated the utility of gene specific markers in identification of unexplored species of related forage crops .

References

- Chandra , A . , Pathak , P . S . , Bhatt , R . K . , 2006 . *Stylosanthes* research in India : Prospects and challenges ahead . *Current Science* 90 , 915-921 .
- Liu , C . J . , Musial , J . M . , 1997 . *Stylosanthes* sp . aff . *S . scabra* : a putative diploid progenitor of *Stylosanthes scabra* (Fabaceae) . *Plant Systematics and Evolution* 208 , 99-105 .
- Phailkaew , C . , Ramesh , C . R . , Kexian , Y . , Stur , W . , 2004 . Utilization of *Stylosanthes* as a forage crop in Asia . In : Chakraborty S . High yielding anthracnose-resistant *Stylosanthes* for agricultural systems . Australia : ACIAR , 65-76 .
- Tewari , S . , Chandra , A . , 2008 . Genetical assessment of diploid progenitors of *S . scabra* by Isozyme , RAPD and STS markers : a possible strategy for improvement of drought tolerant allo-tetraploid *S . scabra* species . *Euphytica* (in press) .