



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

21st International Grassland Congress / 8th
International Rangeland Congress

Genetic Analysis of Genotype X Environment Interaction for Yield Components of *E. canadensis*, *E. sibiricus* and Their Hybrids

Jinghuan Li

Inner Mongolia Agricultural University, China

Jinfeng Yun

Inner Mongolia Agricultural University, China

Shuyan Wang

Inner Mongolia Agricultural University, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/12-1/42>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Genetic analysis of genotype \times environment interaction for yield components of *E. canadensis*, *E. sibiricus* and their hybrids

Li jinghuan^{1,2} Yun jinfeng¹, Wang shuyan¹

¹College of Ecol. and Env. Sci., Inner Mongolia Agric. Univ., Huhhot, Inner Mongolia 010019, P.R. of China; E-mail: lijinghuan0816@126.com

²College of Life Science and Technology, Inner Mongolia Normal University, Hohhot 010022, P.R. of China

Key words : genetic effect, environment interactions, yield components

Introduction *E. canadensis* and *E. sibiricus* are plants whose combining ability and synthesis performance are all good. In this paper, genetic effect of fresh grass yield components of *E. canadensis*, *E. sibiricus* and their hybrids (F_1 and F_2) were studied in two different environments. It was known how the relative size of kinds of genetic diversity and genotype \times environment interactions affect fresh grass yield. All these provided the theory basis for selection of traits in forage breeding for yield and the heterosis utilization of forage.

Materials and methods Single plants of *E. canadensis* (maternal plant), *E. sibiricus* (male parent) and their hybrids (F_1 and F_2) were transplanted at spacing of 60 cm \times 60 cm in the field of experimental farm at DaTai and HaiLiuTu, in Huhhot Inner Mongolia, China, in 2006 respectively. There were 15 plants in each block with three replications in each environment. There were 4 seedlings per hole. During ripening, 5 samples were gotten at random from each block of parents and their hybrids. Quantitative traits, i.e., height, no. of tiller, no. of internodes, stem diameter and yield of grass, were measured with three replication for each sample of parents, F_1 s and F_2 s. All the data were analyzed by QTMModel which was mixed model of genotype \times environment about complex quantitative traits (Zhu J 2004).

Results and discussion Many yield components were affected by not only additive gene action (V_A) and dominant gene action (V_D) but also by genotype \times environment interactions (Table 1). Their importance differed for different yield components, such as no. of tiller, no. of internodes and diameter of stem that were mainly affected by additive gene action (61%, 62% and 44%, respectively), no. of tiller, no. of internode and grass yield were also affected by dominant gene action \times environment interactions (V_{DE}), especially yield of grass (81%). Although plant height was mainly affected by additive gene action \times environment interactions (V_{AE} -56%).

Table 1 Estimated proportions of variance component for yield components.

Parameter	Height	No. of tiller	No. of internode	Stem diameter	Grass yield
V_A/V_P	0.00	0.61*	0.62*	0.44*	0.00
V_D/V_P	0.00	0.00	0.05	0.02	0.12
V_{AE}/V_P	0.13	0.00	0.00	0.01	0.00
V_{DE}/V_P	0.56*	0.03	0.25	0.00	0.81*
V_e/V_P	0.31	0.36	0.08	0.53*	0.07

Conclusions If phenotypic variance of a trait was decided by genetic effects, we may improve the genetic compositions of inbreeding populations by means of genetics and breeding (Zhu, 1992). No. of tiller, no. of internode and diameter of stem can be selected in early generation, but plant height and grass yield were used by tapping Heterosis potential.

References

Zhu J. QGA-Cn[EB/OL]. <http://www.rarlab.com> 2004-01-17/2004-02-01.

Zhu J. Mixed model approaches for estimating genetic variances and covariance (J). *J. Biomath.* 1992, 7(1): 1~11.