



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII
International Rangeland Congress

Summary of Regional Test of Introduced High Quality *Agropyron* Varieties

Nati Jia

Animal Science Academy of Xinjiang, China

Xuanzheng Guo

Animal Science Academy of Xinjiang, China

Li Li

Animal Science Academy of Xinjiang, China

Qiang Cheng

Animal Science Academy of Xinjiang, China

Jie Li

Animal Science Academy of Xinjiang, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/12-2/40>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Summary of regional test of introduced high quality *Agropyron* varieties

Jia Nati¹, Guo xuanzheng¹, Li li¹, Cheng qiang¹, Li jie²

¹Grass Industry Research Institute of Animal Science Academy of Xinjiang, Urumqi, 830000; E-mail: xjndlili@126.com

²Sci. and Tech. Management and International Cooperation Department of Animal Science Academy of Xinjiang, Urumqi, 830000

Key words : variety introduction, forage quality, perennial grasses, regional test

Materials and method Two cultivars of *Agropyron cristatum* cv. Таукумский and Аксенгерский and *Agropyron mongolicum* CK were compared in an experiment located on Desert and Temperate Semi-Arid Steppe in Xinjiang with no-irrigation. The trial design was randomized with three replications. The plot area was 8 8m². Planting was on 20 April 2005, with a row space was 30cm. The experiments were in different ecological areas. Field drought hardiness and physiological ecology were appraised using a 1-5 score (5=drought hardy; 1=drought susceptible). Salt and alkali resistivity was evaluated using 1.0, 1.5, 2.0 and 2.5% NaCl solution to germinate seeds.

Results

Table 1 The growing characteristics of different varieties.

Varieties	Years	Biomass (kg/ha ²)	Regeneration speed (cm/d)	seed yield (kg/ha)	ratio of stem to leaf %
Таукумский	2005-2007	7616	2.18	941.6	69.5
Аксенгерский	2005-2007	8481	2.21	945	69.6
mongolicum(CK)	2005-2007	5637	1.78	734	47.5

Nutrition component For *A. cristatum* Таукумский and аксенгерский respectively the Crude protein (%) were 9.1, 12.2; and NFE (%) 45.5 and 45.3.

Physiological ecology characteristic

Table 2 Physiological ecology characteristic.

Varieties	1.0% NaCl germination(%)	Winter hardiness(%)	Drought hardiness	Resistant lodging	Resistance disease and insect
Таукумский	42	91	+++++	▲▲▲▲▲	resistant
Аксенгерский	44	95	+++++	▲▲▲▲▲	resistant
mongolicum(CK)	0	27	++	▲▲	susceptible

Salt and alkali resistivity When salt concentration of NaCl reached 1.0%, germination percentage of the two introduced varieties of *Agropyron* were 44% and 42%. At higher concentrations of NaCl (1.5, 2.0 and 2.5%), there was no germination.

Reference

Jia nati, Guo xuanzheng, Li jie, The Republic of Kazakhstan *Agropyron* characteristics of germplasm resources [J]. *Pratacultural Science*, 2006, (12): 31-35.