



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## Fertilizer Management for Seed Production of Perennial Forage Crops in the Canadian Great Plains

S. S. Malhi

*Agriculture and Agri-Food Canada, Canada*

C. L. Vera

*Agriculture and Agri-Food Canada, Canada*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/14-1/18>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Fertilizer management for seed production of perennial forage crops in the Canadian Great Plains

S. S. Malhi and C. L. Vera; Agriculture and Agri-Food Canada, P.O. Box 1240, Melfort, Saskatchewan, Canada S0E 1A0; Phone: 306-752-2776 Ext. 230; Fax: 306-752-4911; E-mail: malhis@agr.gc.ca

**Key words** : fertilizer , perennial forages , seed yield

**Introduction** In the Canadian prairies , nitrogen (N) and phosphorus (P) are most commonly deficient nutrients , while potassium (K) and sulphur (S) may also be deficient for certain crop and soil conditions . One or more of these four major nutrients may be limiting for optimum forage seed yield . Perennial forages respond well to the application of fertilizers on nutrient deficient soils . The purpose of this report is to summarize research information on fertilizer management for improving seed production of perennial forages in the Canadian Great Plains .

### Results and discussion

**1) Nitrogen** : For grasses , N fertilizer application produced substantial increase in seed yield , but the magnitude of increase varied with weather conditions in the growing season (Buglass 1964 ; Loeppky et al . 1999) , grass species (Loeppky et al . 1999) and soil characteristics (Thompson and Clark 1989 ; Loeppky et al . 1999) . The best N source for surface-broadcast application was ammonium nitrate (Knowles and Cooke 1952) , but it may not be widely available in the market . Urea is now becoming the most commonly used N fertilizer , but it is highly vulnerable to ammonia volatilization when surface-applied . In general , all N sources can be equally effective when applied properly (e . g . , placed below the soil surface) . Autumn application usually produced higher seed yield than spring application (Knowles and Cooke 1952 ; Lawrence and Kilcher 1964 ; Buglass 1964 ; Ukrainetz 1969) . Legumes coexist with *Rhizobium* or *Sinorhizobium* bacteria in soil that can induce nodulation and convert N<sub>2</sub> from the air to a form that is available to plants . Consequently , no N fertilizer is required for legumes , provided legumes are inoculated with appropriate bacteria (Horton 1991 ; Loeppky et al . 1999) .

**2) Phosphorus** : For grasses , application of P increased seed yield of dryland Altai wildrye grass (Lawrence 1980) , timothy and crested wheatgrass (Loeppky et al . 1999) . Seed yield response of smooth brome grass to N and P fertilizers was also affected by the amount of available N and P in the soil . Unlike N , P is relatively immobile in soil (Malhi et al . 2003) , and tends to have a more lasting residual effect (Lawrence 1980) . For this reason , placement of P fertilizer into the soil , where it will be directly intercepted by roots , is very important (Malhi et al . 2001) . For legumes , application of P can increase longevity of stands by several years , and produced a significant increase in seed yield of alfalfa (Loeppky et al . 1999) .

**3) Potassium** : Improved soil K fertility decreases incidence of winter injury in legume stands by increasing accumulation of carbohydrates in the root system (Bailey 1983) , and increases longevity and seed yield of alfalfa stands . Banding is the preferred method of application .

**4) Sulphur** : Both legume and grasses respond to S fertilizer when soil is deficient in plant-available (S . S . Malhi-unpublished results) .

**5) Micronutrients** : Most soils are adequately supplied with micronutrients and their deficiencies are rare on forages for seed production .

**Conclusions** For grasses , N fertilizer application produced the greatest seed yield increase , when other nutrients were not limiting in the soil , and magnitude of seed yield increase was much higher under cool moist conditions than dry conditions in the growing season . Surface-applied ammonium nitrate was more effective than other ammonium-based fertilizers . Autumn-applied N usually produced higher grass seed yield than spring-applied N . For legumes , focus should be on correcting P , K and S deficiencies , as these plants have the property of fixing N<sub>2</sub> from the air when properly inoculated . Whenever possible , fertilizers (particularly P , K and urea N) should be banded (or injected) into the soil for most efficient use of the nutrients .

### References

- Bailey , L . D . 1983 , Effect of potassium fertilizer and fall harvests on alfalfa grown in the eastern Canadian prairies . *Can . J . Soil Sci .* , 63 , 211-219 .
- Buglass , E . 1964 , Seed production of crested wheatgrass as influenced by various management practices . *Can . J . Plant Sci .* , 44 , 66-74 .
- Horton , R . 1991 , Fertilization for optimum alfalfa seed production in northeastern Saskatchewan . Pages 31-36 in D . W . Goetzen ed . , Proc . 10<sup>th</sup> Annual Canadian Alfalfa Seed Conference , 10-12 January , 1991 , Canadian alfalfa Seed Council , Saskatoon , Saskatchewan , Canada .
- Knowles , R . P . , and Cooke , D . A . 1952 , Response of brome grass to nitrogen fertilizers . *Sci . Agric .* , 32 , 548-554 .