



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

21st International Grassland Congress / 8th  
International Rangeland Congress

---

## An Investigation on Some Ecological Characteristics of *Ziziphus spina chirsti* (L.) in Southern Iran

A. R. Mahmoodi  
*University of Agricultural Sciences, Iran*

Mansour Mesdaghi  
*University of Agricultural Sciences, Iran*

Masoume Amirkhani  
*University of Agricultural Sciences, Iran*

E. Kargari  
*University of Agricultural Sciences, Iran*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/13-2/41>

The 21st International Grassland Congress / 8th International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## An investigation on some ecological characteristics of *Ziziphus spina chirsti* (L.) in southern Iran

A.R. Mahmoodi, M. Mesdaghi, M. Amirkhani and E. Kargari


Department of Range Management, College of Natural Resources, University of Agricultural Sciences, Gorgan, Iran. E-mail: mahmoodi\_150@yahoo.com

**Key words:** *Ziziphus spina chirsti* (L.), distribution, southern Iran

**Introduction** *Ziziphus spina chirsti* is a plant indigenous to the subtropical parts of southern Iran. It is a deciduous shrub growing to the height of 5m. It flowers from July to August and seeds ripen from October to December. This plant has different uses in Iran, China, and Saudi Arabia. The leaves of this plant are locally called "Sedr" and are used for washing the hair and body in Iran. The objectives of this study were to determine phenological and ecological aspects of this species in Iran.

**Material and methods** In this research, the spatial distribution, phenology, and edaphic requirements of *Ziziphus spina chirsti* (L.) were investigated in the Shiraz area by periodical visits to different areas of its distributions. The EC, pH, and texture of soils around this species were measured.

**Results** The general view of *Ziziphus spina chirsti* (L.) and its ecological characteristics are shown in Figure 1 and Table 1.



**Figure 1** The general view of (a) canopy, (b) flowers, and (c) fruits for *Ziziphus spina chirsti* (L.)

**Table 1** Some of the ecological characteristics of *Ziziphus spina chirsti* (L.) in south of Shiraz, Iran.

Distribution	Accompanied Species	Annual rain of study area	Methods of Propagation	Use	Botanical characteristics	Common names	Edaphic requirements
South of Shiraz :	<i>Tecomella</i>			agroforestry			
Larestan	<i>undulata</i>			aforestation	Family of		
Khonj	<i>Mirtous comonis</i>	100-200 mm	seed and stem	medicine	<i>Rhamnacea</i> .	Konar	Loamy texture
Lamerd	<i>Tamarix sp.</i>			fruit	Perennial,	Sedr	EC 2-3.5
				forage	hermaphrodite		pH 7-7.5
				water and soil conservation	flower		

**Conclusion** The economic and medical values of this species are important in south of Iran, so further research is needed for its propagation through seed adaptation in nurseries and areas of its occurrence.

### References

- Duke, J. A., Ayensu, E. S., 1985. Medical plants of China. Reference Publications, Inc. ISBN 0-917256-20-4.
- Ghannadi, A., Tavakoli, N., Mehri-Ardestani, M., 2003. Volatile constituents of the leaves of *Ziziphus spina chirsti* (L.). *Journal of Essential Oil Research*. Bushehr, Iran.
- Zargari, A., 1988. *Medical plants*. University of Tehran publication. 1: 301-303.