



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII  
International Rangeland Congress

---

## Leaf Growth Dynamics and Forage Quality in Temperate and Tropical Grasses

Mónica G. Agnusdei

*Instituto Nacional de Tecnología Agropecuaria, Argentina*

F. Nennin

*Instituto Nacional de Tecnología Agropecuaria, Argentina*

Oscar Norberto Di Marco

*Universidad Nacional de Mar del Plata, Argentina*

M. S. Aello

*Universidad Nacional de Mar del Plata, Argentina*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/15-1/41>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Leaf growth dynamics and forage quality in temperate and tropical grasses


Agnusdei M.G.<sup>1</sup>, Nennin F.<sup>1</sup>, Di Marco O.N.<sup>2</sup>, Aello M.S.<sup>2</sup>. <sup>1</sup>Instituto Nacional de Tecnología Agropecuaria (INTA). <sup>2</sup>Fac. Ciencias Agrarias, Universidad Nacional de Mar del Plata. CC 276. 7620 Balcarce. Argentina. [magnusdei@balcarce.inta.gov.ar](mailto:magnusdei@balcarce.inta.gov.ar)

**Key words** : C<sub>3</sub>/C<sub>4</sub> species , forage quality , indigestible NDF , leaf aging

**Introduction** In grasses , maximum leaf-blade size is achieved at ligulae appearance . However , cell wall (CW) accumulation , as well as the occurrence of changes in CW chemical and structural features continues during leaf aging . Such ontogenic changes in forage quality are currently described by the increase of indigestible NDF (INDF) . At present , many studies are focused on the INDF kinetics during leaf development with modeling purposes . The aim of this study was to evaluate the *in vitro* kinetic of INDF accumulation in individual leaves along their development in three species differing in their potential leaf length and photosynthetic path-way .

**Materials and methods** Studied species : temperate/C<sub>3</sub> : perennial ryegrass (*Lolium perenne* , PRG) ; tropical/C<sub>4</sub> : grama rhodes (*Chloris gayana* , CG) and pangola grass (*Digitaria eriantha* , PG) . The experience was carried out in a glasshouse under adequate nutrient and water availability conditions . Leaf appearance was recorded on marked tillers . Blades were clipped at each leaf appearance interval . The material was immediately frozen with liquid N and further separated into leaf insertion classes . Leaf blade length was measured and leaves of the same insertion level were pooled , lyophilized and ground to 1 mm . INDF was determined in samples incubated in heat sealed F57 filter bags for 24 hours (Daisy , ANKOM Tech . Corp . , Fairport , NY) .

**Results and discussion** INDF accumulation during leaf development was properly described , in the three species , by lineal functions (Figure 1) . This indicates the period of indigestible cell wall formation would have lasted the whole leaf life spans . This pattern differs with the negative exponential saturation curve used in annual ryegrass by Groot and Neuteboom (1997) to describe the evolution of the indigestible cell wall during leaf development . Slopes did not differ between PRG and PG , both being significantly lower than CG (the half , on average) . Average final leaf lengths were 180±16 , 120±10 and 463±19 for PRG , PG and CG . These results stress the influence that leaf length might have on forage quality associated attributes . Moreover , they do not support the idea that forage quality is strictly associated with plant metabolic pathways . The common pattern of INDF increase with leaf aging observed in PRG and PG might , in part , have been associated with the later being shorter leaved than the former at a same leaf age . The obtained results underline the close link existing between plant morphology , leaf aging and forage quality .



**Figure 1** Relationship between indigestible neutral detergent fiber (INDF) and leaf aging (growing degree days :GDD) in C<sub>3</sub> and C<sub>4</sub> perennial grasses .

### Reference

Groot J.J. and J.H. Neuteboom 1997 . Composition and digestibility during ageing of Italian Ryegrass leaves of consecutive insertion levels . *Sci Food Agric* . , 75 , 227-236 .