



University of Kentucky
UKnowledge

International Grassland Congress Proceedings

XXIV International Grassland Congress /
XI International Rangeland Congress

Formononetin of Red Clover (*Trifolium pratense* L.) as Affected by Water Availability

Fernando Ortega-Klose
INIA, Chile

Rafael López-Olivari
INIA, Chile

Muriel Melo
INIA, Chile

Andrés Quiroz
Universidad de La Frontera, Chile

L. Bardehle
Universidad de La Frontera, Chile

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/24/2/33>

The XXIV International Grassland Congress / XI International Rangeland Congress (Sustainable Use of Grassland and Rangeland Resources for Improved Livelihoods) takes place virtually from October 25 through October 29, 2021.

Proceedings edited by the National Organizing Committee of 2021 IGC/IRC Congress

Published by the Kenya Agricultural and Livestock Research Organization

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Formononetin of Red Clover (*Trifolium pratense* L.) as affected by water availability

Ortega-Klose, F. *; López-Olivari, R.¹; Melo, M.¹; Quiroz, A.²; Bardehle, L.³.

* Instituto de Investigaciones Agropecuarias, INIA Carillanca, km 10 camino Cajón-Vilcún s/n, Casilla 929, Temuco, Chile. www.inia.cl Email fortega@inia.cl; ¹Instituto de Investigaciones Agropecuarias, INIA Carillanca; ² Universidad de La Frontera, Laboratorio de Ecología Química, Departamento de Ciencias Químicas y Recursos Naturales, www.ufro.cl Email andres.quiroz@ufro.cl; ³ Universidad de La Frontera, Departamento de Producción Agropecuaria.

Key Words: Formononetin, isoflavones, water availability, plant survival

Abstract

Red clover is a valuable legume in the world. Among the biotic factors limiting persistence of the species in Chile is the root borer *Hylastinus obscurus* (Marsham). Our previous studies have shown a negative effect of formononetin content of the plant on the root borer fitness. The purpose of this research was to assess the relationship between water availability and formononetin concentration. Two independent growth chamber experiments were established at Carillanca Research Center, INIA-Chile in order to study the effect of water availability on formononetin concentration in the aerial and root parts of the plants. In experiment I, three levels of water applied were studied: T1: non-stressed, maintained between 100 and 60% of the readily available soil water (RAW); T2: 60 and 30% of RAW; T3: <30% of RAW. Experiment II consisted of four water levels: T1: 100-80% RAW; T2: 80-60% RAW; T3: 60-40% of RAW; T4: <40% of RAW. In both experiments a complete randomized design was used. For sampling, plants were dug up from the pots and cut at the crown to separate the aerial part from the crown and roots. Formononetin concentration was evaluated by extracting with a methanol solution and relative quantifications based on HPLC-MS. Different water available levels were studied. Formononetin was much higher in the aerial part compared to roots. There were no significant effects of readily available soil water (RAW) over formononetin concentration.

Introduction

Red clover (*Trifolium pratense* L.) is a valuable legume around the world. The main limitation of the species is the lack of persistence related to the high mortality of plants due to a complex of biotic and abiotic factors (López-Olivari and Ortega-Klose 2020). There is a close relationship between plant population and forage yield potential (Ortega et al. 2014). Among the biotic factors associated with red clover mortality in Chile, the main one is the root borer *Hylastinus obscurus* (Marsham). Our previous studies of the relationship plant-borer mediated by semi-chemicals have shown that formononetin is the main isoflavone of red clover eliciting an antifeedant response from *H. obscurus* (Quiroz et al. 2017; Quiroz et al. 2018). The purpose of this research was to study the relationship between soil water availability and formononetin concentration.

Methods and Study Site

Two independent experiments were established at Carillanca Research Center (38°41'SL and 72°25'WL), INIA-Chile, in a growth chamber maintained at 24/18°C day/night (14/10 hours) and 125 Photosynthetically Active Radiation (PAR) during the day. In both experiments, the diploid cultivar Superqueli INIA was sown in greenhouse in seed trays and transplanted later to pots of 12.4 l each (22 cm deep; Andisol type soil with a loam texture). Fertilization was applied according to soil analysis, estimating the doses per pot to an equivalent of 40-113-329 kg/ha of N, K₂O and P₂O₅, respectively. In the experiment I, three levels of water applied were: T1: non-stressed, maintained between 100 and 60% of the readily available soil water (RAW); T2: 60 and 30% of RAW; T3: <30% of RAW. In this experiment I, 5 plants per pot were established with 36 pots in a complete randomized design with three levels of water applied, 2 evaluation dates and 6 pots per level and evaluation date. The experiment II consisted of four water levels: T1: 100-80% RAW; T2: 80-60% RAW; T3: 60-40% of RAW; T4: <40% of RAW. In this experiment II, 9 plants per pot were established with 32 pots in a complete randomized design with four levels of irrigation, 2 evaluation dates and 4 pots per water level and evaluation date. Water was replenishing based

on the weight of each pot. Experiment I was cut and evaluated the first time after 16 days of establishing the water treatments and then evaluated again 33 days after the first cut. Experiment II was evaluated 36 days after establishing water treatments and again 48 days later. For sampling, plants were dug up from the pots and cut at the crown to separate the aerial part from the crown and roots; crown and roots were carefully washed. After sampling, the aerial and root tissues were stabilized immediately by immersion in liquid N₂ for later lyophilization. Yield of the aerial and root sections was evaluated (mg / g DM). Formononetin concentration was evaluated by extracting with a methanol solution and relative quantifications based on HPLC–MS. HPLC analysis was performed using a Diode Array Detector and a C18 reversed-phase column according to the methodology reported by Ramos et al. (2008). Moisture availability per pot (W/W) was evaluated weekly by the weight of each pot. In the experiment II, before each sampling date, the stomatal conductance using a steady-state porometer (SC-1 leaf porometer, METER Group, Inc. USA) was measured. Also, the chlorophyll concentration in the leaf was measured once in the experiment II using a portable chlorophyll meter (SPAD-502, Minolta Camera Co. Ltd., Japan). Data was analyzed by ANOVA and means separated by Duncan (5%).

Results

Figure 1 shows the weight of pots during the second period of experiment II; the four water levels were maintained with three irrigations in this experimental period. There were clear responses in stomatal conductance and chlorophyll content. Stomatal conductance differences were significant and in accordance with water levels (Figure 2). Chlorophyll content was significantly higher with water deficit treatments (T1: 43,9 c; T2: 46,2 c; T3: 52,4 b; T4: 62,4 a). There was no significant response in dry matter yield of both aerial and root sections.

The highest content of formononetin was found in leaves and stems compared to roots. In the experiment I, there were non-significant tendencies to increase formononetin concentration in the aerial part with water stress; something similar occurred in experiment II (Figure 3).

Figure 1. Average weight of pots under four levels of irrigation in the Experiment II. FC: Field capacity; WP: Wilting point. T1: 100-80% of the readily available soil water (RAW); T2: 80-60% RAW; T3: 60-40% RAW; T4: <40% RAW.




Figure 2. Stomatal conductance ($\text{mmol/m}^2/\text{s}^1$) of each treatment before two sampling dates of Experiment II. Different letters within dates indicate significant differences (Duncan's test, $p \leq 0.05$). T1: 100-80% of the readily available soil water (RAW); T2: 80-60% of RAW; T3: 60-40% of RAW; T4: <40% of RAW.






Figure 3. Formononetin content (mg/g) of red clover aerial part (shoots and leaves) in two experiments and two sampling dates each. Experiment I: T1: non stressed, maintained between 100 and 60% of the readily available soil water (RAW); T2: 60 and 30% of RAW; T3: <30% of RAW. Experiment II: T1: 100-80% of RAW; T2: 80-60% of RAW; T3: 60-40% of RAW; T4: <40% of RAW. Bars indicate the standar error of the means.



Experiment I. Sampling 1.



Experiment I. Sampling 2.



Experiment II. Sampling 1



Discussion

times after a period of 3 weeks treatment (De Rijke et al., 2005). By contrary, our results did not show an increase of formononetin content in treatments with water stress; moreover, there was no significant effect of available soil water on formononetin concentration in the aerial part and roots of red clover. Considering the repellent effect elicited by formononetin on *H. obscurus*, the non-effect shown by water availability can be considered a good result since the genetic expression of formononetin content in selected red clover genotypes will be independent of the soil available water level.

Conclusions

There was a higher concentration of formononetin in the aerial parts of red clover compared to roots. There was no significant effect of soil water availability on formononetin content; this is a positive finding if a red clover genetic line with a higher formononetin content is selected since the expression of the trait will be independent of the water soil available level. It is necessary to study other abiotic soil factor (phosphorus level as example) and confirm this findings in field conditions.

Acknowledgments

This work was supported by INIA grant 500302-70, FONDECYT grants 1070270, 1100812,1141245 and 1181697.

References

- Al-Tawaha, A.M., Seguin, P., Smith, D.L. and Bonnell, R.B. 2007. Irrigation level affects isoflavone concentrations of early maturing soya bean cultivars. *J. Agronom. Crop Sci.* 193: 238–246.
- Bennett, J.O., Yu, O., Heatherly, L.G., and Krishnan, H.B. 2004. Accumulation of genistein and daidzein, soybean isoflavones implicated in promoting human health, is significantly elevated by irrigation. *J. Agric. Food Chem.* 52: 7574–7579.
- De Rijke, E., Aardenburg, L., Van Dijk, J., Ariese, F., Ernst, W.H.O., Gooijer, C. and Brinkman, U. 2005. Changed soflavone levels in red clover (*Trifolium pratense* L.) leaves with disturbed root nodulation in response to waterlogging. *J. Chem. Ecol.* 31: 1285-1298.
- Gutierrez-Gonzalez, J.J., Wu, X., Gillman, J.D., Lee, J.D., Zhong, R., Yu, O., Shannon, G., Ellersieck, M., Nguyen, H, and Sleper, D. 2010. Intricate environment-modulated genetic networks control isoflavone accumulation in soybean seeds. *BMC. Plant. Biol.* 10: 105. doi: [10.1186/1471-2229-10-105](https://doi.org/10.1186/1471-2229-10-105).
- López-Olivari, R. and Ortega-Klose, F. 2020. Response of red clover to deficit irrigation: dry matter yield, populations, and irrigation water use efficiency in southern Chile. *Irrig. Sci.* <https://doi.org/10.1007/s00271-020-00693-0>
- Ortega, F., L. Parra and A. Quiroz. 2014. Breeding red clover for improved persistence in Chile: a review. *Crop and Pasture Science* 65: 1138-1146.
- Ortega, F., Quiroz, A.; Parra, L.; Mutis, A.; Hormazábal, E. y Melo, M. 2015. Persistence of Red Clover (*Trifolium pratense* L.) is highly related to plant population: preliminary studies on isoflavones that could act as insect deterrents. Proceedings of the 23th International Grassland Congress, New Delhi, INDIA, Paper ID: 1248.
- Quiroz, A., L. Bardehle, E. Hormazábal, F. Ortega & A. Mutis. 2018. Differential formononetin content in cultivars and experimental lines of red clover (*Trifolium pratense* L.) plants affect the feeding behaviour of *Hylastinus obscurus* (Coleoptera: Curculionidae). *Blacpma* 17(4): 372-380. *Boletín Latinoamericano y del caribe de plantas medicinales y aromáticas*. ISSN 0717 7917
- Quiroz, A., L. Méndez, A. Mutis, E. Hormazábal, F. Ortega, M.A. Birkett & L. Parra. 2017. Antifeedant activity of red clover root isoflavonoids on *Hylastinus obscurus*. *Journal of Soil Science and Plant Nutrition* 17(1): 231-239.
- Ramos, G.P., Dias, P.M.B., Morais, C.B., Froehlich, P.E., Dall’Agnol, M., & Zuanazzi J.A.S. 2008. LC Determination of four isoflavone aglycones in red clover (*Trifolium pratense* L.). *Chromatographia* 67: 125-129.