



University of Kentucky  
UKnowledge

---

International Grassland Congress Proceedings

XXIV International Grassland Congress /  
XI International Rangeland Congress

---

## Sustainable Intensification in Crop-Livestock Systems

P. Rovira  
*INIA, Uruguay*

Paulo C. de F. Carvalho  
*Universidade Federal do Rio Grande do Sul, Brazil*

J. Terra  
*INIA, Uruguay*

F. Lattanzi  
*INIA, Uruguay*

R. Pizzio  
*INTA, Argentina*

*See next page for additional authors*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/24/3/21>

The XXIV International Grassland Congress / XI International Rangeland Congress (Sustainable Use of Grassland and Rangeland Resources for Improved Livelihoods) takes place virtually from October 25 through October 29, 2021.

Proceedings edited by the National Organizing Committee of 2021 IGC/IRC Congress

Published by the Kenya Agricultural and Livestock Research Organization

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

**Presenter Information**

P. Rovira, Paulo C. de F. Carvalho, J. Terra, F. Lattanzi, R. Pizzio, and W. Ayala

# Sustainable intensification in crop-livestock systems

Rovira, P\*; Carvalho, P.C.F\*\*, Terra, J\*, Lattanzi, F\*, Pizzio, R\*\*, Ayala, W.\*

\* Instituto Nacional de Investigación Agropecuaria (INIA, Uruguay); \*\*Universidade Federal do Rio Grande do Sul (UFRGS, Brasil); \*\*\*Instituto Nacional de Tecnología Agropecuaria (INTA, Argentina).

**Key words:** pasture-crops; grasslands; biome; beef cattle

## Abstract

Extensive livestock production is the main animal source food system in the Pampas and Campos sub-regions in South America. Beef cattle and sheep convert forage biomass that humans cannot eat into nutrient-dense human-edible foods (meat and milk) and valuable co-products (wool). However, diverse global pressures are acting on Pampas systems including increasing demand for protein sources (food security), climate change, environmental footprint, and competition for land. The integration of small areas of pasture-crop rotations is an alternative for increasing efficiency and sustainable intensification of agroecosystems based on natural grasslands. Rotations led by improved pastures increase the carrying capacity and productivity of traditional Pampas systems, and include crop production (soybean, rice) as an opportunity for farm diversification and resilience. Therefore, mixed crop-livestock systems can feed more people in terms of calories and protein than what is possible with specialized systems. The higher carrying capacity in the area under the rotation may allow less intensive grazing management in larger areas of native grasslands reducing threats to biodiversity. Some of the integrated management practices in crop-livestock systems are perennial pastures to mitigate soil organic carbon losses during cropping, biological nitrogen fixation by legumes, grazing cover crops, crop residue grazing, dual-purpose crops, and harvesting grain and hay for livestock feed. Also, complementary crop and livestock production systems facilitate more efficient nutrient cycling and self-sufficient systems promoting the circular economy concept as a strategic approach toward system sustainability. Key to understanding the potential of mixed crop-livestock systems are productive, environmental, economic, and social factors that determine their performance, as well as trade-offs among them. Development, implementation, maintenance, and analysis of long-term crop-livestock platforms research provides science-based information to address complex biological systems, and to implement innovative public policies at national scale to regulate soil use and to prevent non-sustainable agricultural practices.

## Introduction

The Pampas and Campos region comprises parts of Argentina and Brazil and the whole of Uruguay (Modernel et al. 2016). It has different subregions defined by vegetation communities, soils, and landscape characteristics (Baldi and Paruelo 2008). The expansion of the agricultural area under continuous cropping and livestock overgrazing of native grasslands are two of the main concerns compromising the sustainability of the Pampa and Campos biome (Medan et al. 2011; Oliveira et al. 2017; Tiscornia et al. 2019). Cultivation of soybean was the most important reason for land use change in the XXI century causing a decline in the area available for livestock. However, cattle numbers remained relatively stable indicating that cattle stocking rates increased in native grasslands areas and decreased in areas dominated by crop production (Modernel et al. 2016). Recoupling livestock and cropping pathways into integrated crop-livestock systems (ICLS) provides an opportunity to achieve sustainable intensification and protect grasslands from conversion to permanent cropland (Smart et al., *in press*) or overgrazed areas. The objective of this short paper is to review key results of the 'Palo a Pique' long term experiment and to discuss the synergy between ICLS and native grasslands in enhancing the sustainability in the Pampa and Campos biome in South America.

## Methods and Study Site

Data shown in the present paper corresponds to a long-term crop-livestock experiment established in 1995 in Treinta y Tres (Uruguay) at the National Institute of Agricultural Research (INIA) 'Palo a Pique' research unit (33°15'54.4"S 54°29'28.1"W). The dominant soils are Typic Argiudols with low to moderate soil fertility (1.5 to 2.0% soil organic carbon content (SOC), mass base, in 20 cm depth), occupying a landscape of gently sloping hills where the erosion risk is moderate to high (Terra et al., 2002). The experiment evaluates four different soil use intensities (rotations) with different combinations of pastures and crops under no-till

technology (Terra et al., 2006; Rovira et al., 2020). The rotations are: (i) continuous cropping without pastures (CC), (ii) short rotation (SR, two years crops and two years of red clover-based pastures), (iii) long rotation (LR, two years crops and four years of white clover-based pastures), (iv) permanent improved white clover-based pasture (PP) without crops. Each phase of the rotations was represented by a paddock of 6 ha, which was the experimental unit (EU), totaling 12 EUs. Although all phases of the rotations were present at the same time, there were no synchronic replications of each phase. Pastures and forage crops are subjected to direct grazing by growing and finishing Aberdeen Angus cattle.

## Results

After 20 years of running the experiment (1995-2015), no soil organic carbon (SOC) differences were found between rotations that included pastures (SR and LR), but they had 21% greater SOC than CC. In addition, SR and LR had lower SOC than PP, while a SOC decreasing trend of 12% was observed in PP compared to the original undisturbed soil under native grasslands located next to the experimental site (Terra and Macedo, 2015). No differences were found in crop yield between the different crop–livestock rotations, averaging ( $\pm$  s.d.)  $2499 \pm 888$ ,  $4871 \pm 1758$ ,  $2771 \pm 1158$ , and  $1569 \pm 479$  kg/ha for soybean, sorghum, wheat, and oat, respectively, across rotations in the period 2005-2016.

From the animal production standpoint, average liveweight gains ( $\pm$  s.d.) for the first four years of the experiment were  $485 \pm 151$  (CC),  $527 \pm 61$  (SR),  $484 \pm 82$  (LR), and  $338 \pm 103$  (PP) kg/ha per year. In the same period, average forage production was 8.5, 10.2, 9.1, and 7.9 t DM/ha per year for CC, SR, LR and PP, respectively, allowing an animal carrying capacity of 951, 726, 722, and 480 kg LW/ha/year for CC, SR, LR, and PP, respectively (Terra and García-Préchac, 2001). During similar period (1992-2004) and in the same research unit, forage production of native grasslands averaged 3.4 t DM/ha per year, with a seasonal distribution of 35%, 26%, 11%, and 28% in summer, autumn, winter, and spring, respectively (Bermudez y Ayala, 2005). The annual production of the year with the highest production (2002) was three times greater than the production of the year with the lowest production (2004) (5.2 and 1.2 t DM/ha, respectively). For an average year, livestock production in native grasslands ranged between 118 and 145 kg LW/ha/year at a stocking rate between 300 and 420 kg LW/ha (Ayala and Bermudez, 2005).

## Discussion

In the Pampas and Campos region the area occupied by grasslands has been reduced by the expansion of soybean. When crops are introduced, rotation with pastures is one of the most basic principles to achieve a sustainable farming system. Here, we describe a large-scale, long-term experiment evaluating four different pasture-crop rotations integrated to livestock production. After 20 years of the experiment, results showed a significant SOC reduction under no-till CC relative to the other rotations containing high proportion of perennial pastures in their cycles. Terra et al. (2006) suggested that the higher content of SOC in pasture-based rotations was related to the greater biomass partitioned to the root systems compared to CC. Recently, evaluating the four rotations using sophisticated models suggested that perennial pastures underpin soil C and nitrogen (N) cycling in crop rotations by maintaining soil C closer to saturation (Pravia et al., 2019). Similar results were obtained in Argentina, where a minimum of 3 years of pastures maintained soil properties within acceptable limits and meet the goals of sustainable intensification in pasture-crop rotations (Studdert et al., 1997).

One of the main features of the experiment is that all paddocks were subjected to direct grazing by beef cattle in a rotationally grazing system adjusting the stocking to pasture conditions. The stocking rate and intensity of grazing determines the amount of residual biomass which is a key parameter for the sustainability of the system. Residual biomass after grazing not only influence animal performance but also represents a physical barrier to compaction through animal trampling, influencing soil aggregation and water dynamics, while avoiding runoff and soil losses from erosion (Carvalho et al. 2018). In Argentina, Fernández et al. (2011) found that the inclusion of livestock to graze cover crops and perennial pastures in ICLS did not produce deterioration of soil physical conditions; although soil penetration resistance was slightly increased within the first 7.5 cm. However, shallow compaction of the topsoil may develop after years of no-till seeding, even in the absence of grazing (Lavado and Taboada, 2009).

Pastures in ICLS increased 2.5 times the forage production and doubled the animal carrying capacity compared to native grasslands. As a result, LW production in pasture-crop rotations more than tripled compared to LW production in native grasslands. Production intensification is one way to reduce carbon footprint per animal and per hectare (Kanter et al., 2016). In Uruguay, Picasso et al. (2014) found that for

every 10 kg increase in productivity (kg LW/ha/year), carbon footprint decreases by 1.2 kg CO<sub>2</sub>e/kg LW and 36 kg CO<sub>2</sub>e/ha. Increasing productivity can lower environmental impacts by spreading baseline emissions in higher animal outputs and by reducing the cattle age at slaughter, which also produces more food on less land area (Sánchez Zubieta et al., 2021). In addition to increase LW production, ICLS provide product and economic diversification through grain harvesting during the cropping phase of the rotations which can be sold or used to feed the animals within the system.

Native grasslands are key partners of ICLS. Coupling strategic and small areas of pasture-crop rotations with larger areas of grasslands increases farm sustainability by reducing the risk of conversion to continuous agriculture (Smart et al., 2020) and overgrazing of grasslands (Lemaire et al., 2015). Temperate pastures composed by grasses (i.e. tall fescue, annual cover crops) and legumes (i.e., lotus, red and white clover) in ICLS supply high-quality forage during winter when production is reduced in grasslands, usually dominated by native C4 species (Modernel et al., 2016). This complementary can alleviate overgrazing of native grasslands usually grazed by animals all year around at a relatively constant stocking rate, although its forage production shows large seasonal variation. On the other hand, livestock can be moved to grasslands when the area under the pasture-crop rotation has limited carrying capacity due to conjunctural (i.e. excessive soil moisture in winter) or structural (i.e. area under fallow or recently seeded in autumn and spring) factors inherent to ICLS. To avoid spread of introduced species from pasture-crop rotations to grasslands areas via cattle feces, it is recommended to have a transient or buffer area where cattle can graze for few days before starting to graze the natural pasture area. The role of native grasslands in ICLS is clearly a topic that merits further research.

In conclusion, strategic areas of pasture-crop rotations integrated to livestock grazing are not the problem but part of the solution to protect vast areas of native grasslands from continuous agriculture and overgrazing. Increasing overall productivity while reducing its environmental footprint and conserving the biodiversity in ICLS represents a great challenge due to specific trade-offs among productive, socioeconomic and environmental goals (Latawiec et al., 2014). This poses a major research challenge in identifying and facilitating sustainable farming systems in the Pampas and Campos region. Research platforms, as the ‘Palo a Pique’ long-term experiment, play a key role to provide scientific data supporting sustainable pathways of intensification.

## Acknowledgements

We thank the Instituto Nacional de Investigación Agropecuaria (INIA) from Uruguay for providing funds to carry out the ‘Palo a Pique’ long-term experiment. We also thank all the past and present field workers, technicians and students that participated in the collection and analysis of data derived from the experiment.

## References

- Ayala, W., Bermúdez, R. 2005. Estrategias de manejo en campos naturales sobre suelos de lomadas de la region Este. In: Seminario de Actualización Técnica en Manejo de Campo Natural (ed. Gómez Miller, R., Albicete, M.M.). Serie Técnica 151: 41-50.
- Baldi, G., Paruelo, J.M. 2008. Land-Use and Land Cover Dynamics in South American Temperate Grasslands. *Ecol. Soc.* 13:6.
- Bermúdez, R., Ayala, W. 2005. Producción de forraje de un campo natural de la zona de lomadas del Este. In: Seminario de Actualización Técnica en Manejo de Campo Natural (ed. Gómez Miller, R., Albicete, M.M.). Serie Técnica 151: 33-40.
- Carvalho, P.C.F., Peterson, C.A., Nunes, P.A.A., Martins, A.P., de Souza Filho, W., Bertolazi, V.T., Kunrath, T.R., Moraes, A., Anghinoni, I. 2018. Animal production and soil characteristics from integrated crop-livestock systems: towards sustainable intensification. *J. Anim. Sci.* 96: 3513-3525.
- Fernández, P.L., Álvarez, C.R., Taboada, M.A. 2011. Assessment of topsoil properties in integrated crop-livestock and continuous cropping systems under zero tillage. *Soil Res.* 49: 143-151
- Kanter, D.R.; Schwoob, M.H.; Baethgen, W.E.; Bervejillo, J.E.; Carriquiry, M.; Dobermann, A.; Ferraro, B.; Lanfranco, B.; Mondelli, M.; Penengo, C.; Saldias, R.; Silva, M.E.; Soares de Lima, J.M. 2016. Translating the sustainable development goals into action: A participatory backcasting approach for developing national agricultural transformation pathways. *Glob. Food Secur.* 10: 71-79.
- Lavado, R.S., M.A. Taboada. 2009. The Argentinean Pampas: A key region with a negative nutrient balance and soil degradation needs better nutrient management and conservation programs to sustain its future viability as a world agresource. *J. Soil Water Conserv.* 64: 150A-153A.
- Lemaire, G., Franzluebbers, A., Carvalho, P.C.F., Dedieu, B. 2014. Integrated crop–livestock systems: Strategies to

- achieve synergy between agricultural production and environmental quality. *Agric., Ecosyst. Environ.* 190: 4-8.
- Medan, D., Torretta, J., Hodara, K., de la Fuente, E., Montaldo, N. 2011. Effects of agriculture expansion and intensification on the vertebrate and invertebrate diversity in the Pampas of Argentina. *Biodivers. Conserv.* 20: 3077-3100.
- Modernel, P., Rossing, W.A.H., Corbeels, M., Dogliotti, S., Picasso, V., Tittonell, P. 2016. Land use change and ecosystem service provision in Pampas and Campos grasslands of southern South America. *Environ. Res. Lett.* 11: 113002.
- Oliveira, T.E., Freitas, D.S., Gianezini, M., Ruviano, C.F., Zago, D., Mercio, T.Z., Dias, E.A., Lampert, V.N., Barcelos, J.O.J. 2017. Agricultural land use change in the Brazilian Pampa Biome: the reduction of natural grasslands. *Land Use Policy* 63: 394-400.
- Pravia, M.V.; Kemanian, A.R.; Terra, J.A.; Shi, Y.; Macedo, I.; Goslee, S. 2019. Soil carbon saturation, productivity, and carbon and nitrogen cycling in pasture-crop rotations. *Agric. Syst.* 171: 13-22.
- Picasso, V.D.; Modernel, P.D.; Becoña, G.; Salvo, L.; Gutiérrez, L.; Astigarraga, L. 2014. Sustainability of meat production beyond carbon footprint: a synthesis of case studies from grazing systems in Uruguay. *Meat Sci.* 98, 346-354.
- Rovira, P., Ayala, W., Terra, J., García-Préchac, F., Harris, P., Lee, M. R. F. Jordana Rivero, M. 2020. The 'Palo a Pique' Long-Term Research Platform: First 25 Years of a Crop–Livestock Experiment in Uruguay. *Agron.* 10: 441.
- Sánchez Zubieta, A., Savian, J.V., Souza Filho, W., Osorio Wallau, M., Marín Gómez, A., Bindelle, J., Bonnet, O.J.F., Faccio Carvalho, P.C.F. Does grazing management provide opportunities to mitigate methane emissions by ruminants in pastoral ecosystems? *Sci. Total Environ.* 754: 142029.
- Smart, A.J., Redfearn, D., Mitchell, R., Wang, T., Zilverberg, C., Bauman, P.J., Derner, J.D., Walker, J., Wright, C. 2020. Forum: Integration of Crop-Livestock Systems: An Opportunity to Protect Grasslands from Conversion to Cropland in the US Great Plains. *Rangel. Ecol. Manag.* (in press, <https://doi.org/10.1016/j.rama.2019.12.007>).
- Studdert, G.A.; Echeverría, H.E.; Casanovas, E.M. 1997. Crop-Pasture Rotation for Sustaining the Quality and Productivity of a Typic Arguidoll. *Soil Sci. Soc. Am. J.* 1997, 61, 1466–1472.
- Terra, J.A.; García-Préchac, F. 2001. Siembra directa y rotaciones forrajeras en las lomadas del este: síntesis 1995-2000. Serie Técnica 125, INIA Treinta y Tres; Treinta y Tres, Uruguay. (In Spanish).
- Terra, J.A.; García-Préchac, F. 2002. Soil organic carbon content of a typic Arguidol in Uruguay under forage crops and pastures for direct grazing: Effect of tillage and rotation system. In Proceedings of the 25th Southern Conservation Tillage Conference for Sustainable Agriculture, Auburn, AL, USA, 24–26 June 2002; pp. 70–74.
- Terra, J.; García-Préchac, F.; Salvo, L.; Hernández, J. 2006. Soil use intensity impacts on total and particulate soil organic matter in no-till pasture-crop rotations under direct grazing. In: Soil Management for Sustainability, 1st ed., Horn, R.; Fleige, H.; Peth, S.; Peng, X., Ed., *Adv. Geoecol.* 38, 233–241, Reiskirchen, Germany.
- Terra, J.A.; Macedo, I. 2015. Twenty years no-till pasture-crop rotation systems impacts on soil organic carbon. Proceedings of the 20th International Soil Tillage Research Organization Conference (ISTRO), September 14-18, 2015 Nanjing, P.R. China.
- Tiscornia, G., Jaurena, M., Baethgen, W. 2019. Drivers, process, and consequences of native grassland degradation: insights from a literature review and a survey in Río de la Plata grasslands. *Agron.* 9: 239.