



Reasons for the Premature Decline in *Astragalus adsurgens* Stands in Kerqin Sandy Land

Q. Zh. Sun

Chinese Academy of Agricultural Sciences, China

Z. L. Wang

Chinese Academy of Agricultural Sciences, China

J. G. Han

China Agricultural University, China

Y. W. Wang

China Agricultural University, China

G. R. Liu

Grassland Service Station of Chifeng City, China

Follow this and additional works at: <https://uknowledge.uky.edu/igc>



Part of the [Agricultural Science Commons](#), [Agronomy and Crop Sciences Commons](#), [Plant Biology Commons](#), [Plant Pathology Commons](#), [Soil Science Commons](#), and the [Weed Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/20/satellitesymposium3/3>

The XX International Grassland Congress took place in Ireland and the UK in June-July 2005.

The main congress took place in Dublin from 26 June to 1 July and was followed by post congress satellite workshops in Aberystwyth, Belfast, Cork, Glasgow and Oxford. The meeting was hosted by the Irish Grassland Association and the British Grassland Society.

Proceedings Editor: D. A. McGilloway

Publisher: Wageningen Academic Publishers, The Netherlands

© Wageningen Academic Publishers, The Netherlands, 2005

The copyright holder has granted the permission for posting the proceedings here.

Reasons for the premature decline in *Astragalus adsurgens* stands in Kerqin sandy land

Q.Zh. Sun¹, Z.L. Wang¹, J.G. Han², Y.W. Wang² and G.R. Liu³

¹Grassland Research Institute, Chinese Academy of Agricultural Sciences, Huhhot, 010010, China, Email: Sunqz@126.com, ²Department of Grassland Science, College of Animal Science and Technology, China Agricultural University, Beijing, 100094, ³Grassland Service Station of Chifeng City, Chifeng, 024000, China

Keywords: *Astragalus adsurgens* cv. Shadawang, degradation, phosphorus, root rot disease

Introduction Diseases partly account for reductions in *Astragalus adsurgens* stand longevity. The effect of some cultural practices on the control of pests and diseases have been reported (Hou, 1986; Nan, 1996), but few reports have detailed the relationship among soil fertiliser status, diseases and premature stand decline. This study was conducted to investigate these relationships in order to extend the longevity of *Astragalus adsurgens* stands.

Materials and methods *Astragalus adsurgens* for this research was established in 1993 at 6-7kg/ha. No irrigation or fertilisers were applied after establishment. Disease incidence was calculated as

$$\text{Disease incidence} = \frac{\text{The number of infected plants} \times 100\%}{\text{The total number of investigated plants}}$$

Plants infected index was divided into 5 classes based on symptoms according to Flood & Isaac (1978) and Latunde-dada & Lucas (1982). Disease index of the stands was calculated according Liu Ruo (1998) as

$$\text{Disease index} = \frac{\sum (\text{Plant infected index} \times \text{plant number}) \times 100\%}{5 \times \text{total number of plants}}$$

Results In the third and fifth year after establishment, phosphorus content in soil, plant leave and stem significantly decreased compared to the first year (Table 1); however, root rot diseases and plant mortality increased. The pathogens accounting for the root rot disease were *Fusarium solani*, *Fusarium oxysporum* and *Fusarium moniliforme*. The stand showed premature degradation in the 5th to 6th year after establishment.

Table 1 The changes of phosphorus in soil and in *Astragalus adsurgens* plants

Years after establishment	Phosphorus content (P ₂ O ₅ mg/ 100g)			Stand condition (%)	
	Soil	Leaf	Stem	Plant mortality	Disease index
0	1.86 a	—	—	—	—
1	1.54 a	0.5401 a	0.0710 a	66.3	54.8
2	—	—	—	78.9	55.6
3	0.54 b	0.1407 b	0.0291 b	100.0	77.7
4	—	—	—	100.0	86.2
5	0.11 c	0.1066 c	0.0233 c	100.0	92.6
6	—	—	—	100.0	95.0

Note: data in a column followed by different letters are significantly different (p<0.05)

Conclusion Premature degradation of *Astragalus adsurgens* were attributed to phosphorus deficiency in soil and root rot diseases caused by the fungi *Fusarium solani*, *F. oxysporum*, and *F. moniliforme*.

References

- Flood J.& I. Issac (1978). Reaction of some cultivars of lucerne to various isolates of *Verticillium albo-atrum*. *Plant Pathology*, 27,166-169.
- Hou, T. J. (1986). Diseases of *Astragalus adsurgens* in western Liaoning province and Inner Mongolia, *Grasslands of China*, 3, 40-43.
- Latunde-dada, A.O. & J.A. Lacas (1982). Variation in resistance to *Verticillium* wilt within seeding populations of some varieties of lucerne. *Plant Pathology*, 31,179-186.
- Liu, R. (1998). Science of Grassland Protection. Agriculture Press of China. 259-263.
- Nan, Zh. B. (1997) *Astragalus adsurgens* disease and their distribution characters. *Acta Prataculturae Sinica*, 14, 30-34.