

**Supplementary Materials for: Latent Profile Analysis of the Five Facet Mindfulness Questionnaire in a Sample with a History of Recurrent Depression**

Table S1.

Chi-square test statistics for pairwise differences between profiles in the test sample ( $N = 343$ ) and validation sample ( $N = 340$ ).

Mindfulness-related variable	Profile comparison	Chi-square test statistic, $p$ -value	
		Test sample	Validation sample
SCS total scale	1 (very low mindfulness) vs. 2 (high mindfulness)	30.25, $p < .001$	86.62, $p < .001$
	1 vs. 3 (moderate mindfulness)	29.38, $p < .001$	38.64, $p < .001$
	1 vs. 4 (non-judgmentally aware)	14.00, $p < .001$	6.90, $p = .009$
	2 vs. 3	8.67, $p = .003$	29.40, $p < .001$
	2 vs. 4	1.15, $p = .283$	23.84, $p < .001$
	3 vs. 4	2.09, $p = .148$	1.43, $p = .233$
SCS self-kindness	1 vs. 2	26.43, $p < .001$	64.78, $p < .001$
	1 vs. 3	26.30, $p < .001$	18.75, $p < .001$
	1 vs. 4	9.98, $p = .002$	0.18, $p = .673$
	2 vs. 3	6.65, $p = .010$	26.29, $p < .001$
	2 vs. 4	2.74, $p = .098$	33.32, $p < .001$
	3 vs. 4	0.30, $p = .581$	6.47, $p = .011$
SCS self-judgement	1 vs. 2	20.20, $p < .001$	62.74, $p < .001$
	1 vs. 3	10.88, $p = .001$	27.29, $p < .001$
	1 vs. 4	11.31, $p = .001$	27.95, $p < .001$
	2 vs. 3	7.27, $p = .007$	19.98, $p < .001$
	2 vs. 4	0.12, $p = .734$	0.83, $p = .361$
	3 vs. 4	3.73, $p = .053$	6.59, $p = .010$
SCS common humanity	1 vs. 2	18.82, $p < .001$	23.34, $p < .001$
	1 vs. 3	12.76, $p < .001$	12.85, $p < .001$
	1 vs. 4	8.17, $p = .004$	0.70, $p = .403$
	2 vs. 3	5.61, $p = .018$	4.25, $p = .039$
	2 vs. 4	0.77, $p = .381$	20.98, $p < .001$
	3 vs. 4	1.25, $p = .263$	13.78, $p < .001$
SCS isolation	1 vs. 2	10.15, $p = .001$	36.30, $p < .001$
	1 vs. 3	16.66, $p < .001$	7.12, $p = .008$
	1 vs. 4	7.90, $p = .005$	18.99, $p < .001$
	2 vs. 3	1.37, $p = .243$	18.19, $p < .001$
	2 vs. 4	0.01, $p = .919$	0.28, $p = .599$
	3 vs. 4	1.32, $p = .251$	8.54, $p = .003$
SCS mindfulness	1 vs. 2	36.11, $p < .001$	41.57, $p < .001$
	1 vs. 3	13.92, $p < .001$	20.07, $p < .001$
	1 vs. 4	6.31, $p = .012$	1.02, $p = .314$
	2 vs. 3	13.72, $p < .001$	10.58, $p = .001$
	2 vs. 4	5.80, $p = .016$	33.69, $p < .001$
	3 vs. 4	0.20, $p = .656$	18.17, $p < .001$
SCS over-identification	1 vs. 2	13.07, $p < .001$	67.61, $p < .001$

	1 vs. 3	12.20, $p < .001$	28.58, $p < .001$
	1 vs. 4	4.75, $p = .029$	9.83, $p = .002$
	2 vs. 3	3.15, $p = .076$	20.71, $p < .001$
	2 vs. 4	0.14, $p = .706$	10.20, $p = .001$
	3 vs. 4	0.80, $p = .373$	0.001, $p = .977$
BDI-II	1 vs. 2	72.81, $p < .001$	47.80, $p < .001$
	1 vs. 3	28.55, $p < .001$	12.93, $p < .001$
	1 vs. 4	38.82, $p < .001$	6.97, $p = .008$
	2 vs. 3	12.51, $p < .001$	24.20, $p < .001$
	2 vs. 4	0.11, $p = .737$	13.20, $p < .001$
	3 vs. 4	5.30, $p = .02$	0.30, $p = .584$
Age of onset of depression	1 vs. 2	1.27, $p = .261$	0.52, $p = .471$
	1 vs. 3	0.87, $p = .352$	0.42, $p = .519$
	1 vs. 4	0.006, $p = .937$	0.08, $p = .781$
	2 vs. 3	0.14, $p = .704$	1.70, $p = .192$
	2 vs. 4	0.69, $p = .405$	0.86, $p = .355$
	3 vs. 4	0.42, $p = .515$	0.13, $p = .718$
Number of previous episodes of depression	1 vs. 2	0.37, $p = .546$	0.60, $p = .438$
	1 vs. 3	2.34, $p = .126$	0.13, $p = .721$
	1 vs. 4	0.16, $p = .693$	0.04, $p = .841$
	2 vs. 3	3.53, $p = .060$	1.00, $p = .318$
	2 vs. 4	0.001, $p = .973$	0.36, $p = .551$
	3 vs. 4	2.11, $p = .146$	0.35, $p = .552$

*Note.* Negatively-phrased items in the SCS (from the self-judgement, isolation, and over identification subscales) were reverse-scored prior to analysis. BDI-II = Beck Depression Inventory; SCS = Self-Compassion Scale.