


Semisynthetic aurones inhibit tubulin polymerization at the colchicine-binding site and repress PC-3 tumor xenografts in nude mice and myc-induced T-ALL in zebrafish

Yanqi Xie^{1,2}, Liliia M. Kril^{1,2}, Tianxin Yu^{1,3}, Wen Zhang^{1,3}, Mykhaylo S. Frasinuk^{1,2,4}, Svitlana P. Bondarenko⁵, Kostyantyn M. Kondratyuk⁴, Elizabeth Hausman¹, Zachary M. Martin^{1,2}, Przemyslaw P. Wyrebek^{1,2}, Xifu Liu⁶, Agripina Deaciuc⁷, Linda P. Dwoskin⁷, Jing Chen¹, Haining Zhu¹, Chang-Guo Zhan^{2,7,8}, Vitaliy M. Sviripa^{2,3,7}, Jessica Blackburn¹, David S. Watt^{1,2,3,7*} & Chunming Liu^{1,3*}

¹Department of Molecular and Cellular Biochemistry, College of Medicine, University of Kentucky, Lexington, KY 40536-0509, USA. ²Center for Pharmaceutical Research and Innovation, College of Pharmacy, University of Kentucky, Lexington, KY 40536-0596, USA ³Lucille Parker Markey Cancer Center, University of Kentucky, Lexington, KY 40536-0093, USA ⁴Institute of Bioorganic Chemistry and Petrochemistry, National Academy of Science of Ukraine, Kyiv 02094, Ukraine ⁵National University of Food Technologies, Kyiv, 01601, Ukraine ⁶Center for Drug Innovation and Discovery, Hebei Normal University, Shijiazhuang, Hebei 050024, People's Republic of China ⁷Department of Pharmaceutical Sciences, College of Pharmacy, University of Kentucky, Lexington, KY 40536-0596, USA ⁸Molecular Modeling and Pharmaceutical Center, College of Pharmacy, University of Kentucky, Lexington, KY 40536-0596, USA. Correspondence and requests for materials should be addressed to CL (email: dwatt@uky.edu; chunming.liu@uky.edu)



Supplementary Figure 1, uncut gel for Figure 3C in manuscript.