

| Table A1. Additional Secondary End Points of the DBT Period                                  |                  |   |   |     |                 |      |
|--|------------------|---|---|-----|-----------------|------|
| Variable   | Placebo (n = 45) |   | Telotristat Ethyl (three times per day) |     |                 |      |
|  | Mean (SD)        | P | 250 mg (n = 45)                         |     | 500 mg (n = 45) |      |
|  | Mean (SD)        | P | Mean (SD)                               | P   | Mean (SD)       | P    |
| <b>Flushing</b>  |                  |   |   |     |                 |      |
| Change from baseline in daily flushing episodes averaged over 12 weeks, counts per day       | -0.16 (1.16)     |   | -0.30 (1.31)                            |     | -0.53 (1.34)    |      |
| Arithmetic mean treatment difference   | —                |   | -0.13                                   |     | -0.36           |      |
| Hodges-Lehmann estimator*  | —                | — | 0.036                                   | .39 | 0.00            | .84  |
| <b>Abdominal pain</b>  |                  |   |   |     |                 |      |
| Change from baseline in abdominal pain averaged over 12 weeks, points†                       | -0.23 (1.16)     |   | -0.49 (1.44)                            |     | -0.33 (1.18)    |      |
| Arithmetic mean treatment difference   | —                |   | -0.26                                   |     | -0.11           |      |
| Hodges-Lehmann estimator*  | —                | — | -0.17                                   | .28 | -0.05           | .87  |
| <b>Daily rescue short-acting SSA use</b>   |                  |   |   |     |                 |      |
| Change from baseline in use of short-acting SSAs, averaged over 12 weeks, injections per day | 0.18             |   | -0.11                                   |     | 0.03            |      |
| Arithmetic mean treatment difference   | —                |   | -0.30                                   |     | -0.15           |      |
| Hodges-Lehmann estimator*  | —                | — | 0                                       | .19 | 0               | .16  |
| <b>Stool consistency</b>   |                  |   |   |     |                 |      |
| Change from baseline in stool consistency averaged over 12 weeks, points                     | -0.22 (0.48)     |   | -0.26 (0.47)                            |     | -0.36 (0.41)    |      |
| Arithmetic mean treatment difference   | —                |   | -0.05                                   |     | -0.15           |      |
| Hodges-Lehmann estimator*  | —                | — | -0.09                                   | .57 | -0.15           | .052 |
| <b>Urgency to defecate</b>   |                  |   |   |     |                 |      |
| Proportion of days   | 0.75 (0.29)      |   | 0.67 (0.34)                             |     | 0.60 (0.31)     |      |
| Arithmetic mean treatment difference   | —                |   | -0.09                                   |     | -0.15           |      |
| Hodges-Lehmann estimator*  | —                | — | -0.02                                   | .35 | -0.13           | .006 |

Abbreviations: DBT, double-blind treatment; SD, standard deviation; SSA, somatostatin analog.  
\*Nonparametric measure derived as the median of all possible differences between the groups.  
†Abdominal pain based on patient rating using an 11-point scale: 0, no pain; 10, worst pain ever experienced.

Table A1. Additional Secondary End Points of the DBT Period